

PCT

WORLD INTELLEC  
In



INTERNATIONAL APPLICATION PUBLISHED

WO 9603508A1

(51) International Patent Classification 6 :

C12N 15/34, 7/01, 15/86, A61K 48/00,  
39/235

A

(43) International Publication Date: 8 February 1996 (08.02.96)

(21) International Application Number: PCT/AU95/00453

(22) International Filing Date: 26 July 1995 (26.07.95)

(30) Priority Data: PM 7101 26 July 1994 (26.07.94) AU

(71) Applicant (for all designated States except US): COMMON-WEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; 407 Royal Parade, Parkville, VIC 3052 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BOTH, Gerald, Wayne [AU/AU]; 4 Leslie Street, North Ryde, NSW 2113 (AU). BOYLE, David, Bernard [AU/AU]; 6 Mary Place, Leopold, VIC 3224 (AU). VRATI, Sudhanshu [IN/IN]; National Institute of Immunology, JNU Campus, New Dehli 110 067 (IN).

(74) Agent: F.B. RICE & CO.; 28A Montague Street, Balmain, NSW 2041 (AU).

(81) Designated States: AU, CA, CN, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published  
With international search report.

(54) Title: DNA ENCODING OVINE ADENOVIRUS (OAV287) AND ITS USE AS A VIRAL VECTOR

(57) Abstract

The invention relates to an isolated DNA molecule comprising the genome of ovine adenovirus OAV287, functionally equivalent DNA molecules or portions thereof. The invention also relates to plasmids and viral vectors including the DNA molecules. The invention also relates to methods for delivering non-adenoviral DNA molecules encoding a polypeptide or polypeptides to animals and in particular to grazing animals.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DNA encoding ovine adenovirus (OAV 287) and its use as a viral vector

Technical Field

The present invention relates to a new full length genomic clone derived from a benign adenovirus (OAV287) isolated from sheep in Australia. The present invention also relates to new viral vectors derived from the benign ovine adenovirus and also relates to the use of these vectors for the delivery and expression of nucleic acid sequences encoding functional RNA molecules or polypeptides to animals.

Background of the Invention

Diseases caused by infectious agents and parasite infestations cause health problems and production losses in domestic animals but for many infectious agents no vaccine exists. Consequently, there are major research efforts worldwide to develop new vaccines which can protect against disease.

While some protective antigens from infectious agents and parasites have been identified, their successful use as vaccines requires the development of systems which can effectively deliver the antigen to the host. A variety of recombinant gene expression vectors derived principally from the pox virus family have been employed as these are generally of low pathogenicity. Expression of the foreign protein following infection by the recombinant viral vector may stimulate a protective immune response in the host.

However, no viral vector has all the attributes desirable for all situations. Some vectors are better suited to particular tasks than others because of their biological properties. For example, it has often proved difficult to stimulate an effective mucosal immune response which can protect against disease. In humans, adenoviruses have been given orally to vaccinate against respiratory disease (1). As this involves protection at mucosal surfaces adenoviruses clearly have potential in

this regard. Human adenovirus vectors have also been used to deliver genes to muscle (2) and other tissues. Although adenoviruses do not generally integrate their DNA into the cellular genome, nevertheless, the DNA persists and long term protein expression is observed. Expression of an appropriate antigen from such cells can generate a systemic immune response which may be protective against the homologous disease causing agent.

Known adenovirus genomes are linear double-stranded DNA molecules which have an inverted terminal repeat sequence (ITR) at each end and a protein covalently bound to the 5'-terminal C residue (3). The genome sequence and structure has now been completely determined for human adenoviruses types 2, 5, 12 and 40 and partially for numerous others, including some animal isolates (see Genbank and EMBL Nucleic Acid databases). Human adenovirus type 2 was the first genome to be sequenced but broadly speaking its genome arrangement is conserved among other characterized adenoviruses i.e. early regions E1-E4 and the structural protein homologues can be recognized in similar locations in the genome. In particular, the E1A/E1B region is located at the left hand end of the genome and region E4 is always located at the right hand end of the genome. Early region E3 is always located between the genes for structural proteins pVIII and fiber, although its size and complexity varies between species e.g. from 3kb with at least 10 open reading frames in human adenoviruses to approximately 0.7kb with only two significant open reading frames in murine adenovirus (4, 5). E3 is a key region for the construction of recombinant viruses as it is non-essential for replication in vitro (6). The late, L region is expressed from the major late promoter, MLP and complex splicing generates families of mRNAs which code for most of the structural viral proteins. Proteins IVa2 and IX appear to have their own promoters.

Although there are some human viral vectors available for medical use there are few animal viral vectors suitable for use in veterinary applications. In order to obtain a more suitable animal viral vector the present inventors have purified an ovine adenovirus (OAV287) isolated from sheep in Western Australia. This ovine adenovirus is serologically related to bovine adenovirus type 7 but is genetically distinct from the bovine adenoviruses and other Australian ovine isolates, as shown by comparisons between the ovine and bovine adenoviruses, based on restriction enzyme profiles (8). The genome arrangement of the virus according to the present invention varies significantly from all other known adenoviruses. The adenoviral DNA molecule of the present invention is suitable for use in viral vectors capable of expressing a variety of polypeptides when used for veterinary applications.

#### Summary of the Invention

According to a first aspect, the present invention consists in an isolated DNA molecule comprising a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 or a functionally equivalent nucleic acid sequence. Preferably, the nucleic acid sequence encoding the genome of the adenovirus is substantially as shown in Figure 1.

In a further preferred embodiment of the first aspect of the present invention, the DNA molecule comprises a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) wherein a portion of the adenoviral genome not essential for the maintenance or viability of the native adenovirus deleted or altered.

In a second aspect, the present invention consists in a DNA molecule including at least a fifteen nucleic acid base sequence being substantially unique to the ovine adenovirus (OAV287) nucleic acid sequence shown in Figure 1. In a preferred embodiment of the second aspect of the

present invention, the at least fifteen nucleic acid base sequence encodes a functional element of ovine adenovirus (OAV287). Preferably, the functional element is selected from the group consisting of promoter, gene, inverted terminal repeat, viral packaging signal and RNA processing signal. The inverted terminal repeat of ovine adenovirus (OAV287) comprises the first 46 nucleic acid bases from the 5' end of each strand of the double stranded DNA genome of the virus.

In a third aspect, the present invention consists in a plasmid including the DNA molecule of the first or second aspects of the present invention. Preferably, the plasmid includes the DNA molecule of the first aspect of the present invention wherein the nucleic acid sequence encoding the adenoviral genome is linked to a nucleic acid sequence encoding an origin of replication and a further nucleic acid encoding a marker. Preferably, the nucleic acid sequence encoding the marker encodes for resistance to an antimicrobial agent. More preferably the antimicrobial agent is ampicillin.

In a further preferred embodiment of the third aspect of the present invention, sequences encoding inverted terminal repeats of the adenovirus are joined.

In a fourth aspect, the present invention consists in a viral vector comprising the DNA molecule of the first aspect of the present invention and at least one nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides.

Preferably, nucleic acid sequence encoding the non-adenoviral polypeptide or polypeptides is derived from bacteria, viruses, parasites or eukaryotes. More preferably, the non-adenoviral polypeptide is rotavirus VP7sc antigen, the parasite polypeptide is *Trichostrongylus colubriformis* 17kD antigen, the *Taenia ovis* 45W antigen or the PM95 antigen from *Lucilia cuprina*.

In another form, the present invention consists in a viral vector comprising the DNA molecule of the first aspect of the present invention and at least one nucleic acid sequence encoding a functional RNA molecule. It will be appreciated by one skilled in the art that a functional RNA molecule can include a messenger RNA molecule, an antisense RNA molecule or a ribozyme.

In a fifth aspect, the present invention consists in a method of delivering a DNA molecule having a nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides to a target cell comprising infecting the target cell with a viral vector according to the fourth aspect of the present invention such that the DNA molecule encoding the polypeptide or polypeptides is expressed and the polypeptide or polypeptides is produced by the target cell.

In a sixth aspect, the present invention consists in a method for delivering a DNA molecule having a nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides to an animal comprising administering to the animal a viral vector according to the fourth aspect of the present invention such that the viral vector infects at least one cell of the animal and the infected cell expresses the DNA molecule encoding the polypeptide or polypeptides and produces the polypeptide or polypeptides. Preferably the animal is a grazing animal and more preferably the grazing animal is a sheep.

In another form, the present invention consists in a method for delivering a DNA molecule having a nucleic acid sequence encoding a functional RNA molecule to an animal comprising administering to the animal a viral vector of the fourth aspect of the present invention having a nucleic acid sequence encoding a functional RNA molecule such that the viral vector infects at least one cell of the animal and the infected cell expresses the DNA

molecule encoding the functional RNA molecule and produces the functional RNA molecule.

As used herein the term "functionally equivalent nucleic acid sequence" is intended to cover minor variations in the ovine adenovirus (OAV287) DNA molecule which, due to degeneracy in the DNA code, does not result in the molecule encoding different viral polypeptides. Further, this term is intended to cover alterations in the DNA code which lead to changes in the encoded polypeptides, but in which such changes do not substantially affect the biological activities of these viral polypeptides.

As used herein the term "functional element" is intended to cover nucleic acid sequences that encode promoters, genes, inverted terminal repeats, viral packaging signals and RNA processing signals. It will be appreciated by one skilled in the art that unique sequences from ovine adenovirus (OAV287) that encode these functional elements may be useful in other systems including plasmids and non-ovine adenoviral vectors.

In order that the nature of the present invention may be more clearly understood preferred forms thereof will be described with reference to the following examples and the accompanying drawings.

#### Brief Description of the Drawings

Figure 1 is the nucleic acid sequence of the OAV287 genome beginning at base 1 of the left-hand ITR.

Figure 2 shows the arrangement of OAV287 genes based on homologies detected with Ad2. Regions with question marks are tentative identifications because of the lack of obvious homology.

Figure 3 indicates the major open reading frames in the proposed E1 region of OAV287. Asterisks show the location of possible initiation codons. A previously unidentified gene (p28kD) which codes for a processed structural protein is encoded on the complementary strand.



Figure 4 shows open reading frames in the region of the OAV287 expected to contain E3. However, E3 is missing as the gap between the pVIII and fiber genes is only 197 nucleotides. The site at which the ApaI/NotI polylinker was later inserted is indicated.

Figure 5 shows the major open reading frames in the probable E3 region of OAV287. Asterisks show the location of potential initiation codons. The SallI site which was modified by end-filling and re-ligation and the alternative site at which a polylinker sequence was later inserted into the genome without loss of infectivity is indicated.

Figure 6 is a scheme describing the construction of a plasmid (pOAV287Cm) containing a full-length clone of the OAV287 genome with pACYC184 sequences inserted in the SallI site. Filled in regions show OAV287 sequences. Cross-hatched sequences are derived from plasmids pUC13 or Bluescribe M13+ (Amp<sup>R</sup>), stippled regions from pSELECT (Tet<sup>R</sup>) and open regions from pACYC184(Cm<sup>R</sup>). Only the key restriction sites used for plasmid construction are indicated.

Figure 7 shows a map of the plasmids pOAV100, pOAV200, pOAV600 and pOAV600S. Arrowheads indicate the ITRs and the approximate location of the major late promoter (MLP). The mutated SallI site and sites at which the ApaI/NotI polylinker sequences were inserted are indicated. Light hatching signifies modified Bluescribe sequences inserted in the KpnI site. Linear, infectious genomes (dark hatching) are released by digestion with KpnI.

Figure 8 shows the results of screening ovine adenoviruses OAV100 and OAV200 rescued by transfection of recombinant plasmids pOAV100 and pOAV200 into CSL503 cells. Portions of the genome spanning (A) the mutated SphI site in OAV100 and (B) the ApaI/EcoRV/NotI polylinker insertion site in OAV200 were amplified by PCR together

with the corresponding regions from wild-type OAV287. The products were digested with SphI (A, lanes 3 & 5) and ApaI, EcoRV or NotI (B, lanes 3-5, and 8-10, respectively). (U) indicates undigested samples.

5        Figure 9 is a map of a plasmid pMT used for the assembly of gene expression cassettes. Fragments containing the OAV287 major late promoter and tripartite leader sequences are linked and precede a multiple cloning site for the insertion of genes of interest. A tandem  
10        polyadenylation signal (AATAAA) follows.

      Figure 10 shows a summary of recombinant viruses which have been rescued from the corresponding infectious plasmids and the gene expression cassettes they carry. Cassettes were inserted into the OAV genome between the  
15        pVIII and fibre genes as indicated.

      Figure 11 shows the expression of (A) the *T. ovis* 45W and *L. cuprina* PM95 antigens in CSL503 cells following infection of these cells with OAV205 and OAV210 viruses, respectively and (B) VP7sc expression in CSL503 and bovine  
20        nasal turbinate cells following infection with virus OAV204. (I) Infected cells (U) Uninfected cells. (M) indicates marker proteins of the sizes shown.

      Figure 12 shows expression of VP7sc in (A) CSL503 cells and (B) rabbit kidney and bovine nasal turbinate  
25        cells following infection with OAV206 virus. (I) Infected cells. (U) uninfected cells. (M) indicates marker proteins of the sizes shown.

#### Description of the Invention

##### METHODS

#### 30        Growth and Purification of OAV287

      The virus, isolated from sheep in 1985, was obtained from R.L. Peet, Animal Health Laboratory, Department of Agriculture, Western Australia. The virus isolate was grown in sheep foetal lung cells (line CSL503) and twice  
35        plaque-purified under solid overlay before stocks were prepared. Virus was purified from CSL503 cells as

described previously (18, 22). DNA was extracted from the virus by digestion with proteinase K (23).

#### Cloning of Genome Fragments

5 Molecular techniques for manipulation, modification and transformation of plasmid DNA which were used in the work described below are described in (9) and similar publications. OAV287 DNA was digested with various restriction endonucleases including BamHI, SphI, SmaI and SalI to deduce the location of these sites (18).

10 The adenovirus genome has a protein covalently linked to each end of the linear dsDNA (24). The BamHI A and D fragments of approximately 8kb and 4kb, respectively, were identified as the terminal genomic fragments because their migration into agarose gels was  
15 dependent on the pre-digestion of viral DNA with proteinase K. The internal BamHI fragments B, C, E and F, estimated at 6.2, 5.1, 3.4 and 1.1kb in size respectively, were separated on an agarose gel, recovered and cloned  
20 into BamHI-digested pUC13 using standard ligation and transformation procedures (9). To clone the terminal BamHI A and D fragments, viral DNA (10µg) was digested with proteinase K (50µg/ml in 10mM Tris/HCL, pH8.0, containing 1mM EDTA and 0.5% SDS) at 65°C for 60min to  
25 remove the terminal protein. The DNA was extracted twice with phenol/chloroform, once with ether and recovered by ethanol precipitation. The 3'ends (of unknown sequence) were then digested exo-nucleolytically with T<sub>4</sub> DNA  
30 polymerase (5 units, Toyobo, Tokyo, Japan) in the presence of dATP (100µM) in buffer containing Tris HCL (50mM), pH8.0, MgCl<sub>2</sub> (7mM), 2-mercaptoethanol (7mM) and BSA (10µg/ml) for 15min at 37°C. The DNA was again purified by phenol extraction and ethanol precipitation described above. To remove the single-stranded terminal regions and  
35 create blunt ends the DNA was digested with 1 unit of mung bean nuclease (Pharmacia, North Ryde, Australia) for 10 min at 37°C in buffer containing Na acetate (30mM), pH4.6,

NaCl (50mM) and ZnCl<sub>2</sub> (1mM) before extraction with phenol/chloroform and recovery by ethanol precipitation. Finally the DNA was digested with BamHI (Pharmacia) and the fragments were separated by electrophoresis in low-melting-point agarose. The BamHI A and D fragments were excised, recovered by NACS column chromatography (BRL, Gaithersburg, Md) and ligated with BamHI/HincII-cut plasmid Bluescribe M13<sup>+</sup> (Stratagene, La Jolla, Ca) prior to transformation into *E. coli* JM109. Positive clones carrying fragments of the expected size were identified, restriction digested and confirmed as correct by nucleotide sequencing and comparison with partial sequence determined directly from genomic DNA. This revealed that three 3'-terminal nucleotides were removed during the cloning procedure.

#### Nucleotide Sequencing of the OAV287 Genome

The complete sequence of the OAV287 genome was determined by sequencing the BamHI fragments A-F using the Sanger method (25) and various kits provided by commercial suppliers. Nested deletions were constructed for the five largest fragments using a double-stranded nested deletion kit (Pharmacia). These were sequenced using standard primers. Based on newly determined sequence other nucleotide primers were synthesised using a DNA synthesizer (ABI, Model 391). In this way both strands of the entire genome and the junctions between the fragments were sequenced.

#### Mutagenesis of the OAV287 genome

For the construction of a full length OAV287 clone and subsequent modification of it to create plasmids such as pOAV200 and pOAV600 certain mutations were required. A relevant portion of the genome was subcloned into Bluescribe (Stratagene, La Jolla, Ca) or a similar plasmid which allowed rescue of single stranded DNA. Later it became possible to use dsDNA for mutagenesis. Oligonucleotides of the desired sequence were synthesized,

phosphorylated and used as primers as described by the manufacturers of Muta-gene Phagemid (Biorad Labs, Ca) or Altered sites II (Promega, Wi) mutagenesis kits. Mutations were generally identified by digestion with the appropriate restriction enzyme or by nucleotide sequencing, or both. Genome fragments containing introduced mutations were subcloned to create larger plasmids such as pOAV200 using appropriate unique restriction sites.

10 Construction of a Full-Length Genomic Clone of OAV287

The terminal BamHI A and D fragments (cloned in Bluescribe M13<sup>+</sup>) were each modified by mutagenesis to add the nucleotides lost during cloning and a KpnI site. The last base of the KpnI site incorporated the C at the 5' end of each genomic ITR sequence. This produced plasmids pAK and pDK (Figure 6).

The left hand approximately 21.5kb of the genome was constructed from the BamHI D and B fragments and the SphI A fragment of approximately 13kb. The genomic BamHI B fragment cloned in pUC13 was modified by mutagenesis (GCATGC to GCATCC) to remove the SphI site at position 8287 producing pUC13B. The modified fragment was released by BamHI digestion and cloned into pDK which had been cut with BamHI and dephosphorylated. Colonies carrying the recombinant plasmid pDBM (Figure 6) were identified by screening with an oligonucleotide which spanned the BamHI B/D junction. The SphI A fragment (approximately 13kb) was cloned into the SphI site of pSELECT (Promega) to form pSESPH. This fragment contains a SmaI site near its left hand end which is common to pDBM. The KpnI/SmaI fragment from pDBM was subcloned into pSESPH which had also been cut with KpnI/SmaI to produce pSELLH, a plasmid based on pSELECT which now contained the left-hand approximately 21.5kb of OAV287 DNA.

35 The right-hand end of the genome was constructed from pAK which contains the right-hand approximately 8.6kb

of the genome and overlaps the SphI A fragment. pAK was cut with SalI and ligated with SalI-cut pACYC184, a plasmid of 4.24kb which contains a gene encoding chloramphenicol (Cm) resistance and an origin for DNA replication, to form a pACm (Figure 6). This plasmid was cut with SphI and KpnI to produce the right-hand genomic fragment incorporating the pACYC184 sequences. This was ligated with the left-hand KpnI/SphI fragment of approximately 21.5kb prepared from pSELLH to produce the final plasmid pOAV287Cm (Figure 6). This plasmid replicates stably in E. coli and therefore removes the need to propagate the virus to obtain genomic DNA for further study. The recombinant genome in plasmid pOAV287Cm differs from the wild-type viral genome by the single point mutation in the SphI site (base 8287), by the presence of pACYC184 sequences in the SalI site and by the addition of a GTAC sequence between the ITRs. However, insertion of pACYC184 sequences in the SalI site disrupts two significant open reading frames whose functions are unknown. If either of the gene products was essential for replication, then pOAV287Cm could not produce infectious virus following transfection. To circumvent this potential problem pOAV287Cm was modified further. First, plasmid Bluescribe M13- (Stratagene, La Jolla, Ca.) was cut with HindIII and end-filled. The linear plasmid was then cut with SmaI, blunt-end ligated and transformed. The resulting plasmid contained an ampicillin resistance gene and origin of replication and lacked SalI and SphI sites but retained a unique KpnI site. This plasmid was cut with KpnI and ligated with KpnI-cut pOAV287Cm. Plasmids which were doubly resistant to ampicillin and chloramphenicol were selected and grown. One of these was cut with SalI to release the pACYC184 sequences, religated and transformed. The resulting plasmid pOAV100 contained the AmpR gene and replication Ori inserted in the KpnI site between the ITR's of the genome (Figure 7). This

plasmid replicated stably in *E. coli* strain JM109 when maintained in the presence ampicillin (200µg/ml). Large quantities of plasmid were grown for transfection studies. Transfection of DNA and Virus rescue

5        To determine whether the recombinant genomic clone was infectious, pOAV100 was cut with KpnI to release the linear viral genome and DNA was transfected into CSL503 sheep foetal lung cells using lipofectamine (GibcoBRL). Solution (A) containing plasmid DNA (2-10µg) and 300µl  
10       EMEM (containing hepes + glutamine), but lacking foetal calf serum (FCS) and solution (B) containing lipofectamine (10µl) + 300µl EMEM (containing hepes + glutamine), but lacking FCS were combined, mixed gently and incubated for 45 minutes at room temperature. Subconfluent CSL503 cells  
15       in a 60mm petri dish were rinsed with 3ml EMEM (plus hepes and glutamine) lacking FCS. EMEM (plus hepes and glutamine) but lacking FCS (2.4ml) was added to the mixture of solutions A and B, mixed gently and added to the rinsed CSL503 cells. Cells were incubated for 5 hours  
20       at 37°C in 5% CO<sub>2</sub>. The incubation medium was changed using complete EMEM plus FCS (10%) and cells were incubated at 37°C in 5% CO<sub>2</sub> until virus plaques or cytopathic effect was visible (7-15 days).

      To confirm that viruses rescued from transfection of  
25       pOAV100 and pOAV200 were derived from those plasmids a portion of the genome of wild-type OAV287, OAV100 and OAV200 viruses was amplified by PCR. For OAV100 a primer pair spanning the region of the mutated SphI site at bases 8287-8292 was used. For OAV200 the primer pair spanned  
30       the insertion site for the ApaI/NotI polylinker between the pVIII and fiber genes. Wild-type OAV287 DNA was amplified as a control in each case. DNA amplified from wild-type OAV287 was cut with SphI whereas the DNA amplified from OAV100 was not (Figure 8A). Similarly  
35       OAV200 DNA was cut with ApaI, EcoRV and NotI whereas

OAV287 DNA was not (Figure 8B). Other viruses were similarly characterised by restriction enzyme digestion. Identification of MLP/TLS elements and Construction of pMT

OAV287 TLS elements were identified as follows and  
5 as described (17). mRNAs present in OAV287-infected  
CSL503 cells were copied into cDNA by reverse  
transcription using primers complementary to the IIIa or  
fiber genes. A primer thought to fall within TLS exon 1  
was then paired with each cDNA primer for PCR. DNA was  
10 successfully amplified, cloned and sequenced. This  
identified TLS exons 2 and 3 (which correspond to bases  
8083-8145 and 8350-8412 of Figure 1, respectively) and the  
3' boundary of TLS exon 1 which occurs at base 5044 of  
Figure 1. A second PCR strategy was then used to obtain  
15 MLP and TLS fragments suitable for assembly into pMT. The  
region in Figure 1 between nucleotides 4861 and 5023,  
thought to contain the MLP was amplified by PCR using a  
plus sense primer which added an ApaI sequence at the 5'  
end and a 3' minus sense primer which introduced an NdeI  
20 site by point mutation at base 5012. Similarly, the TLS  
was amplified using a plus sense primer which introduced  
the NdeI site at base 5012 and a minus sense primer which  
was complementary to bases 8396-8412 and which added a  
HindIII site at the 3' end of the PCR product. The PCR  
25 fragments were digested with ApaI/NdeI and NdeI/HindIII,  
respectively and the fragments were cloned into Bluescript  
SK+ (Stratagene) cut with ApaI/HindIII. The resulting  
plasmid was then digested with HindIII/NotI and a  
synthetic oligonucleotide with HindIII/NotI termini and  
30 the sequence shown in Figure 9 was cloned to produce  
plasmid pMT. Genes of interest were then cloned into  
convenient restriction sites in the NCS. Gene expression  
cassettes were subcloned as ApaI/NotI fragments into  
pOAV200 or rescued into infectious virus.



### Infection of cells and expression of antigens

CSL503 and other cells were infected with viruses at a multiplicity of infection of 20pfu/cell as described previously (21). Infection was allowed to proceed for 24-  
5 60 hr. Cells were then incubated in methionine-free medium in the presence of  $^{35}\text{S}$ -methionine to label newly synthesized proteins. The protein of interest was recovered from cell lysates by immunoprecipitation using a specific antiserum against the expressed protein (21).  
10 Recovered proteins were analysed by polyacrylamide gel electrophoresis and detected by autoradiography or using a phosphorimager (Molecular Dynamics).

### RESULTS

To characterise the genome in molecular terms, BamHI  
15 restriction fragments representing the entire OAV287 genome were cloned into various plasmids and sequenced using methods described in Sambrook (9) and similar publications. Sequences were determined on both strands by using nested sets of deletion mutants together with  
20 synthetic oligonucleotide primers which were synthesized from newly determined sequences.

The viral sequence of 29,544 nucleotides (Figure 1) is considerably shorter (by approximately 6.5kb) than the sequence for human adenoviruses but many genes encoding  
25 structural proteins are identified by their homology with their Ad2 homologues (Figure 2). It is clear, however, that the ovine adenovirus genome shows major structural and sequence variations compared with all other adenoviruses studied to date (Figure 2), in the regions  
30 encoding both structural and non-structural proteins. In particular,

(a) the reading frames tentatively identified as forming the E1A/B regions are named principally on the basis of their location in the genome. Very limited  
35 homology can be detected between the 44.5kD open reading frame (orf) and the large T E1B protein of other

adenoviruses. Homology in the putative E1A region of OAV287 has not so far been detected;

5 (b) in other adenoviruses the E4 region is normally located at the right-hand end of the genome. The OAV287 E4? region is tentatively identified based only on the presence of a protein sequence motif HCHC..PGSLQC which is found in 18.8kD and 30.85kD orfs in this region. Identical or very similar motifs are found in the E4 34kD protein of human Ad2 and Ad40 and mouse adenoviruses;

10 (c) the distance between the end of pVIII and the beginning of fiber, which in other viruses defines the E3 region, is only 197 nucleotides (Figure 4). The E3 region equivalent, if it exists in ovine adenovirus, may consist of the cluster of open reading frames which are present in the right to left orientation on the complementary DNA strand, at the right-hand end of the genome (Figures 2 and 15 5). However, these sequences show no detectable homology with any other adenovirus and the functions of these proteins cannot be deduced from such comparisons;

20 (d) there is a region of approximately 1kb which lies between E3? and E4? which has a very high A/T content (70.2%) (Figure 1). As there are no open reading frames encoding greater than approximately 30 amino acids in length on either DNA strand it is unlikely that the region codes for any proteins, unless mRNAs are generated by very 25 complex splicing events. This region has no known equivalent in any other adenovirus;

(e) other differences are apparent in the structural proteins of the virus. OAV287 lacks homologues of Ad2 30 proteins V and IX. However, OAV287 has a completely new gene coding for p28kD which is located on the complementary strand of the E1A? region (Figure 2 and 3). This is a structural protein with an apparent size of 28kD by SDS PAGE which, according to N-terminal sequencing 35 data, is cleaved from a larger precursor. No homology

between this protein and others in the databases has been detected;

(f) in most other genomes the VA RNA genes are located between the Terminal protein and the 52/55k genes. In OAV287 there is no room for them as the reading frames overlap.

These differences serve to emphasize the unique character of the OAV287 isolate compared with other human and animal adenoviruses. In addition, since the OAV287 non-structural regions show little or no homology with equivalent regions in other adenoviruses, sequence comparisons do not reveal the identity of likely non-essential regions of the genome. Moreover the viral DNA cannot easily be manipulated to test for dispensable sequences.

The present inventors have produced a plasmid containing a full length infectious copy of an ovine adenovirus genome in which the ITR sequences are linked by a short sequence which creates a unique restriction enzyme site. A plasmid containing a full length infectious copy of an ovine adenovirus genome linked to a bacterial origin for DNA replication and a marker gene has been produced. Partial clones of OAV287 genomic DNA were specifically modified and initially linked to a gene encoding antibiotic resistance and origin of replication inserted into the unique SalI site of the genome (Figure 6 and see Methods). Such a plasmid can be grown in bacteria and more easily manipulated.

The circular genomic clone differs from the naturally occurring circles that occur in Ad5-infected cells (10) and that might exist in OAV287-infected cells in that the 40 base pair ITRs are joined by a GTAC linker. Together with the last and first nucleotides of the genome (G and C, respectively, see Figure 1), this sequence forms a unique KpnI site (GGTACC) when the ITRs are joined head to tail. Other sites such as EcoRI, BamHI, SalI, KsaI etc

which have recognition sequences beginning with G and ending with C are suitable if they are unique as the 3' and 5' terminal nucleotides of other adenovirus genomes are G and C, respectively. A plasmid with a suitable antibiotic resistance gene e.g. ampR and origin of replication can be inserted at the unique site or elsewhere in the genome to form a plasmid which can be propagated in bacteria. Plasmids propagated in the presence of 200 µg/ml ampicillin in *E.coli* strains JM109 and DH5-alpha retain the KpnI sites and inserted sequences, indicating that the OAV287 ITR sequences are stable when linked in this manner. This approach may therefore be used to engineer other adenovirus genomes. If desired the GTAC linker sequence can be removed and the authentic termini regenerated prior to transfection by digestion with KpnI (or another appropriate enzyme) and incubation with T4 DNA polymerase to create blunt ends (9).

A method for generating linear infectious genomes from circular plasmids involved digesting the circular plasmid containing the full length copy of the OAV287 genome with restriction enzyme KpnI to generate a genome with the authentic 5' nucleotide dCMP. The linear DNA is then introduced into CSL503 cells using lipofectamine as the transfecting reagent.

To develop a viral genome as a vector it is essential to identify region(s) of the genome which are non-essential for function. These regions can be then substituted or deleted to make room for foreign DNA (11, 12), or they may be the site for insertion of foreign DNA. In the human adenovirus genome DNA has been substituted or inserted into the E1 and E3 regions (13, 14, 15) and at the extreme right-hand end of the genome between E4 and ITR, usually with the concomitant deletion of non-essential regions to facilitate packaging of the genome (16). Adenoviruses will package genomes up to ~6% larger

than the wild-type, probably due to physical constraints dictated by the capsid structure (11).

Non-essential sites in the OAV287 genome were identified by insertion of a polylinker sequence  
5 containing *ApaI* and *NotI* restriction sites. This linker was introduced into the genome copy in pOAV100 between nucleotides 22,139 and 22,130 of Figure 1 by site directed mutagenesis to create plasmid pOAV200 (Figure 7). This corresponds to a site located in the intergenic region  
10 between genes for the pVIII and fiber proteins which was chosen because it avoids disruption of RNA processing signals in the region. A transcription termination site for the L4 family of RNAs maps 26 nucleotides upstream and the splice junction between  
15 the tripartite leader sequences and fiber mRNA maps 144 nucleotides downstream of the insertion site, respectively (17). Transfection of pOAV200 into CSL503 cells resulted in the rescue of virus OAV200. The second site at which the polylinker was inserted was located between bases  
20 26,645 and 26,646 of Figure 1. This created plasmid pOAV600 (Figure 7). This insertion site corresponds to the right hand end of the A/T-rich region (Figure 2) whose function and precise boundaries are unknown. The site was chosen as it is six nucleotides to the left of the  
25 transcription termination point for RNAs transcribed from right to left from the E3' region (Figure 2). This was determined by sequencing cloned RT-PCR-amplified cDNAs derived from the region using methods similar to those described for the pVIII/fiber region (17). Transfection  
30 of pOAV600 into CSL503 cells yielded virus OAV600.

The above insertion strategy identified two regions of the genome which can be interrupted and created sites for subcloning gene expression cassettes.

A further non-essential site was identified using  
35 the unique *SalI* site located at bases 28644-28649 of Figure 1. The site was cut with *SalI*, end-filled and

religated to disrupt the reading frames which spanned the site. A plasmid pOAV600S (Figure 7), which had lost the site was identified by digestion with Sall. When pOAV600S was transfected into CSL503 cells, virus OAV600S was recovered. The loss of the Sall site in this virus was confirmed by digesting the viral genome with Sall. As the Sall site falls within two significant open reading frames (which extend on the complementary strand between bases 28457 and 29014 and between 28511 and 28699), which were disrupted by end-filling and religation, the gene products derived from the reading frames are probably also dispensable. This group of reading frames may therefore constitute the E3 region of OAV287 as no other gene products in any adenovirus are dispensable for replication, in vitro. This implies that it should be possible to delete the whole region labelled as E3? in Figure 2. In addition, in other experiments a 1kb NdeI fragment was deleted from the region marked as E4? in Figure 2. This deletion disrupted several reading frames in the region. No virus has been rescued from a such a plasmid, suggesting that it is not dispensable and accordingly, it may be E4.

Many viruses replicate incompletely in heterologous hosts, often entering cells but being unable to produce mature virus particles because of a block in the replication cycle. In the context of recombinant viral vectors, this represents a desirable safety feature, provided that replication is not blocked before appropriate and effective expression of the foreign gene occurs. OAV287 does not replicate productively in heterologous cell types (18), the only exception so far being bovine nasal turbinate cells in which viral titres are significantly reduced compared with the CSL503 cells. Recombinant forms of OAV287 have been constructed to determine whether expression of a reporter gene under the control of an appropriate promoter occurs.

Foreign gene expression requires that the gene be functionally linked to a promoter. This may be a viral promoter inherent in the genome, or a foreign promoter subcloned together with the gene of interest into a suitable site. The promoter driving gene expression must function in CSL503 and preferably a range of other cell types. In this work an OAV287 genomic promoter was used initially. Subsequently an heterologous promoter was also used. In adenoviruses, expression of the structural proteins is driven by the major late promoter (MLP). Families of RNA transcripts derived from the MLP contain a common sequence element, the tripartite leader sequence (TLS) at their 5' ends. The present inventors have identified those nucleotides in the OAV287 genome which comprise the TLS by using RT-PCR amplification of late mRNA transcripts present in OAV287-infected cells and sequencing of cloned cDNAs (17). A candidate MLP was expected to be present just to the left of TLS exon 1 (Figure 2). The MLP and TLS elements were subcloned using PCR techniques into a separate plasmid pMT (Figure 9) and linked with genes of interest. These promoter/gene cassettes were subcloned as ApaI/NotI fragments into the polylinker ApaI/NotI sites of pOAV200. Using this strategy plasmids pOAV203, pOAV204, pOAV205 and pOAV210 were constructed. These incorporate genes encoding a 17kD soluble protein from *T. colubriformis*, a rotavirus VP7sc gene (19), the 45W antigen from *Taenia ovis* (20) and a membrane protein (PM95) from *Lucilia cuprina*, respectively. Plasmid pOAV202, contained the 17kD antigen but lacked the MLP/TLS elements. These plasmids were transfected into CSL503 cells and rescued as viruses OAV202, OAV203, OAV204, OAV205 and OAV210, respectively (Figure 10).

The human cytomegalovirus immediate early IE94 promoter plus enhancer, which functions in a range of human and animal cell types (21), was also linked to the

rotavirus VP7sc antigen gene. This cassette was assembled by replacing the MLP/TLS elements in pMT/VP7sc with the HCMV enhancer-promoter region. The cassette was inserted in pOAV200 to create pOAV206. pOAV206 was transfected  
5 into CSL503 cells and virus OAV206 was rescued (Figure 10).

CSL503 and other cells were infected with the viruses described above and at various times post-infection the cells were radiolabelled with <sup>35</sup>S-methionine. Proteins of interest were recovered from cell  
10 lysates by immunoprecipitation using an appropriate antiserum. Recovered proteins were analysed by polyacrylamide gel electrophoresis and detected by autoradiography.

When virus OAV202 was used, no expression of the *T. coulbriformis* 17kD antigen was observed by immunofluorescence. As this virus lacks the MLP/TLS elements and carries only the 17kD gene this result demonstrates that there is no viral promoter upstream or  
20 adjacent to the insertion point between the pVIII and fiber genes which is capable of driving gene expression. As the E3 region is also missing from this site there is no requirement for a nearby promoter. This situation contrasts with observations made using a human Ad5 E3  
25 recombinant (21). In this case a promoterless gene inserted 3' proximal to the pVIII gene was expressed, probably from the adjacent E3 promoter or the upstream MLP (15, 21). This result further emphasizes the unique  
30 nature of the OAV287 genome. Recombinant OAV287 viruses carrying the MLP/TLS elements were tested for expression in CSL503 cells. With OAV204, expression was easily detected in infected, but not in uninfected cells at 24hr post-infection (Figure 11A). Similarly, when viruses  
OAV205, and OAV210 were tested, gene products of 24kD and  
35 approximately 95kD, respectively were detected (Figure 11B). Therefore it is clear that MLP/TLS elements contain



the necessary information to drive gene expression in the homologous cell line under replication-permissive conditions. However, when OAV204 was tested in a heterologous rabbit kidney cell line in which the virus does not replicate productively, no VP7sc expression was observed. Some replication occurs in bovine nasal turbinate cells, although to a lower titre than in CSL503 cells. In the latter cells, expression of VP7sc was detected following infection with OAV204 (Figure 11B).

Virus OAV206 containing the HCMV enhancer/promoter element linked to the VP7sc gene was used to examine the function of a heterologous promoter in the context of the OAV287 genome. CSL503 cells infected with this virus readily expressed VP7sc antigen at 24-48hr post infection (Figure 12A). With this virus VP7sc expression was also observed in the non-permissive rabbit kidney cell line and in bovine nasal turbinate cells (Figure 12B). These results suggest that the HCMV or a similar constitutive promoter may be preferred over the MLP to drive gene expression in OAV recombinants in non-permissive cells.

One recombinant virus was also administered to sheep. Five sheep were vaccinated intraconjunctivally and intranasally with  $0.7 \times 10^8$  pfu of OAV203. At three days post-inoculation virus was recovered from the nasal swab of one sheep and from the conjunctival swabs of two sheep and confirmed as the recombinant virus by PCR analysis. Animals showed no obvious ill effects from such vaccination.

The viral vectors of the present invention can be used for the delivery and expression of therapeutic genes in grazing animals. In species which are not normally infected by ovine adenoviruses the lack of pre-existing immunity should allow efficient infection, gene delivery and expression. The genes may encode vaccine antigens, molecules which promote growth in production animals, molecules which modify production traits by manipulating

hormone responses and other biologically active or therapeutic molecules. The virus does not replicate productively in many non-ovine cells but the use of heterologous promoters allows the delivery and expression of genes while minimising the possibility of virus spread to a non-target host. As the DNA of adenovirus vectors can persist in cells in an unintegrated form, with the appropriate choice of promoter, expression over a prolonged period can be achieved.

10 References

1. R. B. Couch, et al., *Am Rev Resp Dis* 88, 394-403 (1963).
2. T. Ragot, et al., *Nature* 361, 647-650 (1993).
3. M. S. Horwitz, in *Virology* B. N. Fields, D. M. Knipe, Eds. (Raven Press, New York, 1990) pp. 1679-1721.
- 15 4. W. S. M. Wold, L. R. Gooding, *Virology* 184, 1-8 (1991).
5. K. S. Raviprakash, A. Grunhaus, M. A. El Kholy, M. S. Horwitz, *J Virol* 63, 5455-5458 (1989).
- 20 6. A. M. Lewis, et al., *J Virol* 11, 655-664 (1973).
7. R. L. Peet, W. Coackley, D. A. Purcell, C. W. Robartson, B. M. Micke, *Aust Vet J* 60, 307-308 (1983).
8. M. Benko, A. Bartha, G. Wadell, *Interviol* 29, 346-350 (1988).
- 25 9. J. Sambrook, E. F. Fritsch, T. Maniatis, Eds., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989).
10. F. L. Graham, *EMBO J* 3, 2917-2922 (1984).
11. F. L. Graham, L. Prevec, in *Methods in Molecular Biology* E. J. Murray, J. M. Walker, Eds. (Humana Press, Clifton, NJ, 1991), vol. 7, pp. 1-19.
- 30 12. F. L. Graham, L. Prevec, in *Vaccines: New approaches to immunological problems* R. W. Ellis, Ed. (Butterworth-Heinemann, Stoneham, Ma, 1992) pp. 363-390.
- 35 13. M. Levrero, et al., *GENE (Amst)* 101, 195-202 (1991).

14. J. E. Morin, et al., *Proc. Natl. Acad. Sci. USA* 84, 4626-4630 (1987).
15. G. W. Both, et al., *Virology* 193, 940-950 (1993).
16. P. K. Chanda, et al., *Virology* 175, 535-547 (1990).
- 5 17. S. V. Vratil, D. B. Boyle, R. Kockerhans, G. W. Both, *Virology* 209, In press, (1995).
18. D. B. Boyle, et al., *Vet Microbiol* 41, 281-291 (1994).
19. M. E. Andrew, et al., *J Virol* 64, 4776-4783 (1990).
- 10 20. K. S. Johnson, et al., *Nature* 338, 585-587 (1989).
21. Z. Z. Xu, V. Krougliak, L. Prevec, F. L. Graham, G. W. Both, *J gen Virol* In Press, (1995).
22. S. H. Larsen, H. D., *Virology* 82, 182-195 (1977).
23. E. Nakano, D. Panicali, E. Paoletti, *Proc Natl Acad Sci USA* 79, 1593-1596 (1982).
- 15 24. D. M. K. Rekosh, W. C. Russell, A. J. D. Bellet, A. J. Robinson, *Cell* 11, 283-295 (1977).
25. F. Sanger, S. Nicklen, A. R. Coulson, *Proc Natl Acad Sci USA* 74, 5463-5467 (1977).
- 20 It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all aspects as
- 25 illustrative and non-restrictive.

## CLAIMS:

1. An isolated DNA molecule comprising a nucleotide sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 or a functionally equivalent nucleic acid sequence.
2. The DNA molecule as claimed in claim 1 such that the nucleic acid sequence encoding the genome of the ovine adenovirus is substantially as shown in Figure 1.
3. An isolated DNA molecule comprising a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 wherein a portion of the adenoviral genome not essential for the maintenance or viability of the native adenovirus is deleted or altered.
4. An isolated DNA molecule comprising at least a 15 nucleic acid base sequence being substantially unique to the ovine adenovirus (OAV287) nucleic acid sequence as shown in Figure 1.
5. The DNA molecule as claimed in claim 4 such that the at least 15 nucleic acid base sequence encodes a functional element of ovine adenovirus (OAV287).
6. The DNA molecule as claimed in claim 5 such that the functional element is selected from the group consisting of promoter, gene, inverted terminal repeat, viral packaging signal and RNA processing signal.
7. The DNA molecule as claimed in claim 6 such that the functional element is the inverted terminal repeat having the nucleic acid base sequence 1 to 46 as shown in Figure 1.
8. A plasmid including the DNA molecule as claimed in any one of claims 1 to 7.
9. A plasmid including the DNA molecule as claimed in any one of claims 1 to 3 such that the nucleic acid sequence encoding the adenovirus genome or a portion thereof is linked to a nucleic acid sequence encoding an origin of replication and a further nucleic acid sequence encoding a marker.

10. The plasmid as claimed in claim 9 such that nucleic acid sequences encoding inverted terminal repeats of the adenovirus are joined.
11. The plasmid as claimed in claim 9 or 10 such that  
5 the nucleic acid sequence encoding the marker encodes for resistance to an antimicrobial agent.
12. A viral vector comprising a DNA molecule including a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 or  
10 a functionally equivalent nucleic acid sequence or a portion thereof and at least one nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides.
13. The viral vector as claimed in claim 12 such that the nucleic acid sequence encoding the genome of the  
15 adenovirus is substantially as shown in Figure 1.
14. A viral vector comprising a DNA molecule including a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 wherein a portion of the adenoviral genome not essential  
20 for the maintenance or viability of the native adenovirus is deleted or altered, and at least one nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides.
15. The viral vector as claimed in any one of claims 12 to 14 such that the nucleic acid sequence encoding the polypeptide or polypeptides encodes a polypeptide or polypeptides derived from bacteria, viruses, parasites or eukaryotes.
16. The viral vector as claimed in claim 15 such that  
30 non-adenoviral polypeptide is rotavirus VP7sc antigen, the parasite polypeptide is *Trichostrongylus colubriformis* 17kD antigen, the *Taenia ovis* 45W antigen or the PM95 antigen from *Lucilia cuprina*.
17. A method of delivering a DNA molecule having a  
35 nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides to a target cell, the method

comprising infecting the target cell with a viral vector as claimed in any one of claims 12 to 16 such that the DNA molecule encoding the polypeptide or polypeptides is expressed and the polypeptide or polypeptides is produced by the target cell.

18. A method for delivering a DNA molecule having a nucleic acid sequence encoding a non-adenoviral polypeptide or polypeptides to an animal, the method comprising administering to the animal a viral vector as claimed in any one of claims 12 to 16, such that the viral vector infects at least one cell of the animal and the infected cell expresses the DNA molecule encoding the polypeptide or polypeptides and produces the polypeptide or polypeptides.

19. The method as claimed in claim 18 such that the animal is a grazing animal.

20. The method as claimed in claim 19 such that the grazing animal is a sheep.

21. A viral vector comprising a DNA molecule including a nucleic acid sequence encoding the genome of ovine adenovirus (OAV287) substantially as shown in Figure 1 or a functionally equivalent nucleic acid sequence or a portion thereof and at least one nucleic acid sequence encoding a functional RNA molecule.

22. The viral vector as claimed in claim 21 such that the functional RNA molecule is an antisense RNA molecule or ribozyme.

23. A method for delivering a DNA molecule having a nucleic acid sequence encoding a functional RNA molecule to an animal, the method comprising administering to the animal a viral vector as claimed in claim 21 or 22, such that the viral vector infects at least one cell of the animal and the infected cell expresses the DNA molecule encoding the functional RNA molecule and produces the RNA molecule.

1/23

Fig 1

CTATTCATAT	ATATAACGTT	GCACAGAGGC	GGGGCGTGTG	GGTTTTTTAT	TGTTTATTGT	60
CATGGAATTT	ACAAAGAAGT	AAGTTGTTGG	ATCTTTATTC	ACAATTCCTT	TAACAATGAC	120
TTTTTTACTT	ATTACATTTT	TCATCTTTTT	TACTTCACAT	GATATTTTAC	TTAAATTTTG	180
TACATACAAG	CCAAAATTCG	CATAAAATGT	CTTACTTTAA	AAAGTTAAAT	TTTTTTTTTA	240
ACGCATAAAT	GGACGTACAG	CAGCAATTGG	AATAGCAGGA	AGGGCCATTG	TAAAGTGTGT	300
TCCTGCTGAT	GCCGCTGCAG	AAAGGATAGA	TGCTATCGTA	CGCATAAACC	CCCCTCCTAT	360
TTGTTTCATCT	GCTGCTTTTA	TTATATCTTC	TGCCAATCTA	GGTGATATTT	GCTTTTGAAT	420
GCTGTTTCCA	AAAGCTTGCA	TCATCGGATT	TTCAATTAAA	TGGATTGGAT	TTGCAGAATT	480
TCCTTAAAAA	TAGCCCAACC	CATCTAAAGC	AGTTAAAAAGT	ATTCTCCCTC	CAGGAACCAC	540
AGATATAAAT	AAGCGGAGCA	ACCGAGAGGT	TAAATTCCAG	GGTCCTCCGA	AGAGAGTATC	600
TAGGATCAGG	CCAAGAAAGT	AACCAAAAAG	ACTTGTAAGT	AGAAGTTGTC	TGATATGCTT	660
TGGAGAGGAC	TGTTAAAAAT	GCAAAACGGT	ATCTAATGAC	CATTTCTTCT	TTACTTTTAC	720
ATCTGTATCA	TGTTCTCCAT	CAGAAGGTCT	TATTGGGAAG	TACCATTGGT	CACGAGCATC	780
TTTGAAGACT	TCTGTTTCTT	GAAATTCCTG	TTTCGGTAAG	CGACTAGCAG	TTATGGTATT	840
AGGAATATTG	ACGGTAATGT	TATTCACATC	TACAATTTCT	GGAGGAATCC	ATCTTGCATA	900
GGATGAAATG	GGTTTTGTGG	GTTCTTTCAA	TATATAATTG	CGAGGAGGGT	TTTCCAAAAA	960
TCTCTGAACA	TAAGTATTTT	CTGATTTTGG	CGGTTTTTTG	CTTTTTCGCG	CTCTTTTTCT	1020
TGGCTTTGGT	CTTTGAAAT	TTTTCTTCCT	TTTTCTGTAG	GCTCCTCCTG	CTAAAGCTGT	1080
GTTATTTGTG	ACGTACATCC	TGTTAGCTAC	ACGATTTTCC	CGGACTGCAA	ATTTTTTTGC	1140
CAAATGGAAA	AGAAATTGCT	GAAACCTTCT	ATTAATCATA	TAAATTGTCA	GTGGAATCAT	1200
GAATCAGATA	GTGCAGGATT	TTTTCTTTTT	GATACTGATA	ATTTATACTA	TTATGTATTG	1260
GATCAAGTGT	CTTGGATATG	TTTAAGAGAT	ATAACTCTTC	ATTGTGATCG	CATGTGGTTA	1320
GCGGTTTGT	TTTGTGTTG	CAAATCTAAA	TTTGATGTAC	ACAATATTCT	AGCGGGAGTA	1380
CATGTTATGT	AATGAAAAATG	ACGTCGGGGA	TTGAATGGAT	TGAGCCTTAT	TTGACATTTT	1440
TCTGTGATTT	TTTTGCCTTA	TTAGGAAATA	AATTTGTGGC	GCCAGTACGA	TGGAGATTGG	1500
AATGACTCCT	GCATTTACAG	AAAGGAATTT	GTACTGTGTT	TTGCTTGACT	TTAATTTAAG	1560
ATGGTATCAG	CAGATATTTA	ACCCAATATG	GATTAAGCCA	AATTTATGGG	CTTCTCTGA	1620
TTTTTTAAAA	AAAATGGCCT	TTATTTATGC	TAGCGACTTG	GCGTTGTTAA	ATTCTTACAT	1680
CCCTGGTAAT	GTTTGTAACA	AACTTGATAT	CATCAAGAAA	GATCTTCCTG	AAGATTTTAC	1740
CGTGTCTATG	TTTTGTGTCT	TAGTGTGTTG	GCTTGCTTCT	TTCTGTAAAG	GTTCTAATTT	1800
AGCTGAAACT	CGCCAGAAAT	GTCACGCGGT	AAGCAAATTT	CTGGCACAAC	TATCAAAATT	1860
AATAAAACCC	TAATTTTITAG	TTTGTA AAAA	TAGAATTCAA	ATTTTTAACG	CCACAATGAC	1920
TTCCGGCGGAG	TTTTCTGTTG	AATTTCTTTA	TGTTTCTAAG	CCAATTGTTT	CATGGCCTGC	1980
TTCCGCATCT	TCTAATAATT	CATCGAGTCA	GAATATTGAC	TTTCCTGTTT	TTAAACCAGA	2040
TCAAGATCCA	ATAGCCTTCT	TTCAAACTAA	CAATACGGCT	TACTTACAAC	CTGGAGCTAC	2100
TTATTACTGG	AAGTGTATCG	AACTGTCAAA	GCCTATTAC	ATTTACGGTC	AAGGAGCTAC	2160
AGTACAACCT	GTCGGACCTG	GACCTGTGTT	TGTTTTCAAC	AGTGAAAGTG	TTATTCCTGA	2220
AGATTTTTTAC	GTCGTGTTTG	AAAATATCAA	CTTTATTGAA	GATGAATTTT	CTATTAGAAG	2280
TGGCCAGTTA	AGTTTAGGAC	TTACAACCTCA	CAGTGCTGTA	TGGTTTATCA	ATGTATGGAA	2340
AACTTCAATA	GTCAATTGTA	ACTTTAAAAA	TTTTAGGGGA	GCGGCTCTTT	GGTATTCAGA	2400
TAATAGAAAT	TTTTGGAATG	CGAGAAAAATG	GAATCAGCAG	CATTTAGTTT	CAAATTGTCG	2460
TTTTAATGGT	TGTAGAATTG	GAATTTCTAA	TACTGGTTCA	TCTGAATATT	CCATAGCCAG	2520
TCAAAATCAA	TTTTATGATT	GTCAAATCTG	TTTTAATGTA	ACCGGGGGTA	ATTGGTCTAG	2580
AAATAATAAT	GTTATTGTTA	ACTGTAGATG	TGCTTATCTG	CATGTTGGAG	ATAACATGTG	2640
GTATGAAGGC	CATTCCGAAA	ATAATAATCC	CGCTAAGGGT	ACTTTCTGCA	ATAACATAAT	2700
TAACCATGCT	GATAACGGAG	GCAATGTCTG	GCCTACTCAG	TTTAAACTTA	CAGATGGATC	2760
AACGATACAG	TTAGCATCAT	TTTATTTTGA	TGATAATCAA	GAAATTCCAC	CTTGTTATAG	2820
CGGTAATTTT	CATTGGTTTTG	GAGATGTAAA	CATTGTAAAT	TTTTCTACCA	CAAAAATTGA	2880
TAAATGGTGC	ATTACTGGAT	GTAATTTCTA	TGGTAATACA	CATGCAGCTA	ACGATGCTGG	2940
TCAAGTTCAG	GTTGCTGAAG	CTGTAAAAGA	CAAAGTGTIT	ATTATTGGGT	GTTCTGGTAA	3000
TAATGTAAACC	ATGAAAAATA	TTGTAGAAGG	TAACATGACT	CCAAAAATTG	GTACAATAAA	3060
GTA AAAA ACT	TTTTATTCAA	AACAAAATGG	ATTTACATTT	AAACGTTTTA	CATATTGATT	3120
CTGCGTATAA	GTTCTTTTTT	TAAACACTCT	TCTAATTTCC	ATACATGCTT	GATAAAACAA	3180

2/23

Fig 1 (cont)

ACTTTGTAAA	TTCATAAATA	TAGGTTTGAC	TTGATCAGAA	GGTGAATAAT	AGCTCCATCT	3240
AAATGATTCC	GTAATAGGAA	CATTATTATA	TATTAACCAG	CTATATTTTG	AGTTAACTCT	3300
TGCATGATCC	ACTATATCTT	TAAGTACAGG	GATAAGTGCA	CTCGGAAATC	CAAAAAGAATA	3360
GTTTTTAATA	AATCTATTTA	TCTGTGAAGA	ATCAAGCTGC	GGACTAATAA	CATGACATTT	3420
TGATTGAATT	TTTAAATCCT	TAATATTTCC	TCTATCATGA	CGCGGGTTCA	TATTATGTAA	3480
AACTACTACA	ACAGTGTAAC	CATTACATTT	GGCAAATCTA	TTAAAAATTT	TTGACGGTAA	3540
AGCATGAAAG	AAAGAACTTA	TAGAATGACA	TGATCCCAAT	TGATTCATAC	ATTCATCTAT	3600
TATAATACAG	ATAGATCCTT	CACTTGCAGC	TCTGCAGAAT	ATATTATCTG	GATTATCAAT	3660
ATTTAGATTA	GTATCGGAAA	TAGCATCTTT	GAAAGCTAAT	TGTATAAATT	TTGGATTTAA	3720
TGTTTTTGTT	AGTGGATTAG	AGAATGCATC	GTAGTTTCTT	TCAACACACT	GTGCTTTCCA	3780
CGCAATTTTT	TCTTCTAATG	GAACAGTACC	TTTTTCTGGA	GTTATGAAAA	AAATTGTTTC	3840
TGGTATTGGA	TCAATTAGTT	TTCCAGATAT	AATATTTCTT	ATAAATTGAG	ATTTTCCGCT	3900
ACCTGTGGGT	CCATATACAG	TAACAATGAA	TGGTTGTAAT	CCGCAGTTTA	AACTGGGTAT	3960
ACAGCCATCT	TTTAACAGAT	TGTGAGCCTC	ATTTACAGTT	TTTTGATAAT	TTACAGCAT	4020
ATTGTGTAAA	TCAGTCATAA	GTTGACCATG	ATACATACAT	TTATCAAAAA	CTTCTTGACT	4080
TTCTGGAAAT	GGATTTCTGC	AAATAGAAGG	ATCTATCTTT	ACAACATCAT	TTTTCCAATT	4140
TAATGTGTCA	CTTAAAAATT	TTCCCAAAAA	GGATTTCTG	TCAATGGTTC	TTGCGGTCTT	4200
GGATTTGGGT	GTCTCTTGTC	GTACGGGTAA	AGTAAGTATC	CTTTCTTCCA	CTGGATCCCT	4260
TTCTCATCG	TTTGATCCTT	CCAAGGTCTC	AGAATTCTGG	TTAGTTGCTT	CTCTACCACC	4320
GTGAATGGTA	CATCGGTTCC	ACTTGCAGTT	TGCAGTGTCT	TTTTTAACT	TTCTCTCGAT	4380
GTCTGAAACT	CTTCTGTGG	TTGTTCTAAT	AAATTATAGT	CAGTAAAAACA	ATGTTTTAGA	4440
ATTTCAATAGT	TTAAACAAAT	TTTAGCATGA	CCTTTGGCTC	TTAATTTTCC	TTCTCCAATA	4500
AATTTACAGT	TTTTACAAGT	TATGTCTTTT	AAAGCATATA	ATTTAGGAGC	TAAAATACAT	4560
GTTTCTGAAC	TGAATGCTTC	AGCTCCGCAA	CGGTTACAAA	CAGTTTCGCA	TTCAACCAAC	4620
CAAGTTAGAC	ATGGATGTTT	TTTCATCAAG	ATTAATTTTG	AGTTATATTT	TTTAAGTCTA	4680
TGTAATCCTT	TTGATAACAT	GAGTTGGTGG	CCCTTTTCTG	TTAAGAATAA	CGAGTCTGTA	4740
TCACCATAAA	TACTTTTTAT	CTCCCTTTCT	ATGTAAGGTT	TACCCATATC	TTCCCATAT	4800
AAAAATTTCTG	CCCACCTCAT	CATGAAAGCT	CTGGTCCAAG	CCAGCACAAA	GGATGCTATC	4860
TGAGTTGGAT	ATCGGTTGTT	CTTGATCCAT	TCTTCCTTAT	CCTCAATAGT	TGTTAAAAAT	4920
AAATCATTAC	AATCAGCAGA	TAAAAAAGTT	ATAGGCTTAA	AAGTCACGTG	ATCTTGATTT	4980
CCTATAAAAA	GTGGAAAAAT	AAAATTTTCA	TTTGTGTCTT	TGGAATCTTT	GGGCGGCATT	5040
TCAGGTAGGT	TTGAAAAATA	CTGATTCCAC	TCAAATGAAC	GTTTTGGTAA	TGATTTACTA	5100
ATCAGAGTTG	TGTATGATGT	AATTTACAGT	GATCCATTTT	CTAATCTTTT	TTTATCTTTT	5160
TCTTCAATAT	TTTCAGCAAA	CACACTTTTC	TTTTTATCTA	TACGGGTAGC	AAACGAACCA	5220
TATAAAGCAT	TTGATAACAA	TTTACTTATA	CTTCGCTGAA	TCTTGTGTGT	ACTTTTACTT	5280
GCTTTTCTT	TAGCCATAAT	ATTTACTTTT	ACATATTTT	GACATAACGG	TTTCCAGTCA	5340
CTCCATACAG	CATACATTTT	AGAGCTTTTG	ATTATTTTGC	ATTTCCATCC	TCTATTGTGT	5400
AAGGTGATTA	AATCGATAGA	GGTCAGTACT	TCATTTATCA	ATGTTTCATT	TGACCAGCAT	5460
AACTTTCCAC	TTTTTTTAGA	ACATAATGGA	GGTAACACAT	CAAGATAATC	TAATGATGGG	5520
GGTTCACAA	CGGCTACCAC	AATCATAGGT	TTGATTGAAT	TGTCAAATA	ATCTATTTT	5580
TCTTTTCTTT	GTAGTAGTTC	TTGAAAGTAA	TCTATTTGTG	CATTGGCTTC	AAAAGCATTT	5640
AAAGTTTTTC	CATATGGAAG	TGGATGCGTT	AAGGCACTAG	CATACATTCC	GCAGATATCA	5700
TACACATATA	TTGCTTCTTC	AAATATTCTT	AAAAATGAAG	GATAACATCT	TCCTCCTCTT	5760
AAACTCATTC	TAACAAAATC	ATACATTTTT	TCTGATGGAG	CTTCCAAATT	TCTTAGGAAT	5820
TCAGAGGGAT	GATCTTCTTC	ATTATAAAAG	ATTGTTTAA	ACAATGCTTG	AGTATTACTA	5880
CTAATTGTAG	GACGTTGGAA	TATATTAATA	GAACACTCAA	GCTTTAAAGA	TGTTGTACAG	5940
AACTCTTGAT	AACCTTCTAT	AAGTTTTTCA	ACTAATTGAG	CCGTAACAT	AACATCATCA	6000
ATACAATACT	CCTTAGCTTC	CTCTAATAAG	TTGTATTTT	GGTTGTGTTT	TGGTTTGT	6060
TGTAATATTT	CTTCAATGA	ATTCCAATAT	TTTTGAACTG	GATAACCAT	GTTTTCTTTT	6120
TCATATTCTC	CCAACATAAA	AAAATCATTG	ATTGCCCTGT	AAGGACAATA	ACCTTTGCTA	6180
ACACTCAACT	GATATGCAGT	AGCAGCGTCT	CTTAAAGAAG	AGTGGGTTAA	CAAAAATGTA	6240
TCCCTAACCA	TAAATTTTAT	ACCTTGCCAT	TTCATATCTT	CAAAAATTAAT	AATTCCATTT	6300
TTCCATCTTT	CATAAGTTGT	ATGTGAAGGT	TTCTTAAAGC	AAGGATTTGG	AAGAGATAAT	6360



3/23

Fig 1 (cont)

GTAATATCAT	TAAATAACAG	TTTTCCAGCA	CGAGGCATAA	AGCTTCTTGT	CAGCTTAAAC	6420
ATTGAAAGTT	CTTCACTGTC	TATTCCTTCT	AATACATGAC	TTGCAAGTAT	GATTTTCATCA	6480
AAACCACAGA	TATTATGACC	TACTACATAT	AATTCAATAT	ATCTTGGTTC	GCACTGTTTT	6540
AATTTTTTTT	CITTATTTAA	GACCATGATG	TCTTCATATG	ATAAATTGA	TTCAAGACCA	6600
TGATTTTCAC	AAAACGTTGA	CCAGTATTTT	TAGCTACTG	AAATTGTAG	CTCTGTTCTG	6660
AATTTTTTAA	AAGCTATGCC	AATTTTCATCT	TCTTTTTTAT	TTAACATTAC	AAAACATTCT	6720
CTGTTTACCT	CATAACCTAT	ATCGGTAGCT	ATTTTAGAAG	CAATTTTTAT	GAGTGATTTA	6780
CATCCAATTA	ACTTAAAAAC	CAACAAGTAA	GGAGTAACT	GTTTTCCATA	CAAAGAATGG	6840
TAAGTATATG	TTTCAATATC	ATAAACAATA	AAAAGACGTT	TTGCTTTTAT	GGCTCCAAC	6900
GGATTAAATT	TGATTTTTTC	CCACCAGAGT	TTGTTCAT	GGTGAATATT	GTGATAATAG	6960
AAGTCCCGTC	TTCTGGATGA	GCAGTTGTGT	ATATTACTAT	AAATTGTTCC	GCAGAATTCA	7020
CATTTATTCT	GTTGTTTAA	AGTTTTTATT	AAATATATTT	CTCCTTTTAA	AATCAATAAT	7080
TCTATTGGTA	ACAAATTTCC	ATTAAGAATT	TCTTCAGTCA	TCTTAAAAAA	TCTTTGTGTG	7140
AACTTCCATA	TTTTTAAAGA	TACGGGGGTG	TTAGAAATCAC	AAAGTTTTTAA	AACATCTAAA	7200
ACATTTTCTA	CTTCTTTGAA	AGAATTTAAT	TTTAAACCCCT	GAATTGCAAA	GTAATTATAA	7260
AAACTTTTTT	CAAAATTCTT	GATGATATA	ATTTTATAT	ATGTATCCTC	ATATATTCCA	7320
GTAATATAAG	TAGTAGTTCT	TTGCTTTATT	ATTGCTTTG	AAGCCATCTG	TTTAAAGCCG	7380
CTTCCCGTAC	TCGCTCAAAG	CTTCTTAAAA	CAACTTCATT	TGTACTATAG	CCAACAATTC	7440
CAGACAATTT	TATTTCTAAAT	GCTATTTCAA	CTGAATCTAA	ATCTGAAAAA	TCCGTGTTTA	7500
CTTGGTTGAT	TACTTCTTCT	ATGCTCCAC	TGTCTTCTAC	GAAGTCTATA	TCTTGAAGTA	7560
ATTGGTCTCT	TTCTTCTGGA	GTTGAAAAAG	AGTAAGATCT	TTCATTAGCT	TCTATAATTC	7620
CTAAAAATC	ACGAGTTATT	CTGCTATATA	GTTGCTGAA	TGCTTGTGTT	TCTCTATTAA	7680
ACCAAATCT	AGTAAATATA	TCTTCTCCAT	TTTCATTCT	ACCTCTTAAT	ATAATTTGAA	7740
CAAATTGGAT	TCCAATATTT	CTGGCAGCTA	ACCTATTTTG	CACTAAATTT	AAGTATAAGT	7800
AATATAGCGT	GCTTGCCACA	TGCTCTAATA	TAAAGAAATA	CACTAACCAT	TTTTGAATAA	7860
AATCATCAGT	CAATCTATTT	TCATTATAAA	ATCTAATAAG	TAATTGAAAA	AATTCACCTC	7920
CGTAATTAAT	AAAATTACTC	CTTCTTGCTT	CAGGAGTTAA	TTCTTCTTCT	AAATTTTGAA	7980
TTAAATCTAC	TATTGAAGCT	ATCACTTCAT	CATTAAATTC	TTCCCTACTC	AGATCGCTTG	8040
AGCTCGGCTC	GCGATCTGAA	AATCCTTCAT	CTTCTATTTT	AGGAACAGTA	AGAGGAGAAC	8100
TAGAAGTTTC	TTCAACATTC	CTTACCCTTT	GGCGTCTATT	AACAGGTAAT	CTATCAATAA	8160
ATCTTCTGAT	TACATCACCC	CTTGAACGTC	CTGTATTTT	AGTAATAGCT	CTATAATTTT	8220
CCCTAGGTCT	TAATCTGAAT	GGTAATCCTA	CTCTTGCTCC	TGACCTTAAA	GTTAATGCTC	8280
CACCATGCAT	CCCACCTTTT	CCTAAAGTTA	ATACAGTTGC	TAAATCTTTT	AAATTAATTC	8340
GATTTTCAGC	TTCTGGAATT	TCCAGCTGTG	AAAATTCATC	TATAAAAAAGC	TCAATCCAGA	8400
ATTGAGAAAA	AGGTAAGTCT	AATATACATT	CACTATTATG	CATGTTAGAC	AAAATTTAAA	8460
ATTTACATAA	AGCTTTTTTA	ATTTTACAAA	TTAACTTTAT	AAGGTAAGTA	TCCCTTTCTT	8520
GCAAATTTAA	AACCATAAAA	GCTTGAGAAA	AAGGTTGATA	ATGCTGCTGA	AAAGATCTAT	8580
TCTGATTTTG	AGCTGAAATA	GCGGAGCCAA	AACCTTGCAAT	GTCTGCAAGT	TGCAGACTCC	8640
CTAATATTCT	ATCCATTAAA	ACCGCGTTTT	GAATTTGACT	AATTGTTTGT	GAAAAATTTT	8700
CTACATTTTG	AATTGCTCTC	ATATATGACC	CAGTATTTAT	GGAGTATGAA	CAATCAGTTA	8760
AAATTTGCCA	GGTCATGCGT	CTCTCAAAAC	TTATAGGTGA	AAGATACAAC	TTATATGAAA	8820
TGTTGCTGTA	AGTCCGCTGA	TCAAACAGAT	ACTGGTTTAA	AACTCGCGCC	ACATAAAAAAT	8880
ACCCAATTAA	TAAATTTGGT	GGAGGTTCTC	CTTCAAATGG	TGGTTGTGAA	GTAACAGGTC	8940
CTCTTGGGCG	TAAATCGAGT	AATTGAGTCA	CTGGATAATT	AAAAAATCGA	TTAGCCCAT	9000
TTATTCCCTT	TTCATGTATA	GTCCTTGACC	TGGCAATACT	TCGATTATTA	AGGTCAAGTG	9060
TTAAACGTAA	ATATCGTAAG	GTATGTTGAC	TTTGCCAGT	GAGTTGTTGC	CATTGGTGAA	9120
TCTGCAAGGC	AAACAAAAAA	TTTATCTTAT	TACTGCAGAT	GCATCCTATT	TTACAAAAAT	9180
TACGTTTCATC	ATTGGAAACT	CCAGACTTAT	CAAGCAACTC	CCCGGGCAGC	TCAAATAAAA	9240
ATGAAAAAGA	TGAATTTGAA	CCAGCAGTTG	GCATTTCTAG	CAAACCATCT	GATGAATTTA	9300
ATATGAGACG	ATCTCAAAGA	GATGATAATT	TACCTAAAAAG	TCAGATACCA	GATGATAGATA	9360
TACTACATGA	TAAAAATCCT	AAAATGGCAG	AAGAACGAGA	CTTAATGTAT	AAATCTTCTG	9420
CTTGCAATAA	ACTTGATGAT	TCTAAACAAT	TAAAAACTGA	TATGTTTCAGG	CCGGATTTTG	9480
CTGGAACATG	TCCAGCTCAA	AGACACATAG	AAGCCGCAGA	GCTAAAGAGA	AATGGATCTT	9540

4/23

Fig 1 (cont)

ATACTCGTAG	TTTAGAACAA	TGGACACATG	ATTCTTTTAT	AAGTCATGTT	AAACAATTAC	9600
TTTCTAGACC	ATTTATATCT	CTAGGTATTA	CATATTTGGA	TGATTTTTTG	CAGACTTATT	9660
TAGATCATAC	TGAATCGTCT	TCTTTAAACT	TTCAACTGTT	TACTTTAATA	AATCACTGTT	9720
CAGAAAATAC	TTTAAAACGG	ATTTTAAAC	ACATTTCTAA	AAAAAATGAA	AAAAATCAAT	9780
ATGTAAATCA	ATGGTTGATT	GATCTCATT	CATGTATATA	TCTAATTATA	AGAGATGAAC	9840
AAAATGTTAC	AGAACAAGTT	AATGCCCTTT	TAGTAACTAG	TAATCACTTA	GCTTTACATT	9900
TTGCAAAGAA	AGCTACAGGT	GGATTCTATC	CTACAGCAGA	CAAGTTAGCG	AAGACTCATA	9960
TTTTTTTCAA	GAGAATAATT	TTAGGAATAC	TTTCGCTAGC	AGAAAGTATA	GGTTGCTATA	10020
CTGTGAATCC	ATATTGCAAA	AATCCTTTGA	AAAAGTCAAA	AGTAGAAGTA	GAACCAAGTG	10080
ACGAAATGTA	TATGTTTCAGC	TTAAAAGGTG	CACCTGAACA	TCCTGATTCC	GACGAAGACG	10140
AAGACAGTGG	ACTTCAAAAT	GAATAATTAT	CATAAATGGA	CTTCTAATGT	TATAGATGCA	10200
ATTCTATCAA	ACAAAGCTCT	TTTAGCTATA	AAAATTTTAA	AAGTCAACCG	TTTGCAAACA	10260
AATTGAATGC	TTTGAATCA	GCAGTTGTGC	CTCCAAGAAA	AGATGATACT	CCTGAAATGA	10320
TAGCAAAATCT	TTTAAAAGAA	TTAGTTGCTT	TGGGAGCTAT	TCGCAGTGAT	GAAGTTGGCC	10380
CATTATATTC	TGACCTTCTT	ATCAGAGTTC	ACAAATATAA	TAGCTTGAAT	GTTCAATCAA	10440
ATTTGCAAAAC	TTTAAACAGGA	GACATTAAAT	CACCTCAATC	CGATATAATT	AGAAGTTCCG	10500
ATATTCCCAA	TTTAAAGTAAT	CAAGTTGTTT	TAAATACATT	TTTAAATTCT	TTGCCCTCAA	10560
CTGTTACATT	TGGACAACAT	AATTATGAAG	CTTTTAAACA	AACTCTAAGA	TTATTTGTTA	10620
ATGAGACACC	TAATATTACA	GTTTTTAGAT	CAGGAAATGA	TACTTTAATT	CAGGTTAACA	10680
TAACAGGAAT	TCATACAATT	AATTTGAATG	ATGCATTTAA	AAATTTAAAA	AATTTTTGGG	10740
GAATAGTATT	AACAGGTGAA	TTTATTCCAG	GTGATATTAC	AAGCAGACTA	ACAGCTAATA	10800
CAAGAGTACT	GCTTTATTTT	CTTGCTCCTT	TTACAAATGA	TAATACATTC	ACACCTGATA	10860
CTTTTCTAGC	TTTACTCATG	AAATTATATA	GATTGACAGT	TTCTTCTGCT	TTAGATTTTG	10920
AAGAAGAAAC	TGAAGCTGAA	GTAGAAAATG	TAGCTCAACA	AATAGGATCC	ACTAGTGCAG	10980
ATTTTACAAA	GACTTTAGGA	TATCTATTAA	AAAACAAAGA	AGAATCATTT	TCGCCTCCCA	11040
AATCATTATC	TCCTAGACAA	CTGGGTATTT	TAAGGTTTAT	ACAGAAAAGT	CTGGTAGATA	11100
AAATTGATAG	AAATAATGAA	GATCCATGGG	ATGCTTTAGA	AACTTTATCT	TATTCATTTT	11160
CTCCGTCATT	TTATGAGGCC	AATGGGCCCT	TTATTAGACG	GTAAATAACT	TATATGGAAT	11220
TTGCCCTTACG	TAATTCTCCT	ACTTACTTCA	GAGAAATTTA	CTCCAACAAA	TATTGGATAC	11280
CACCCAATTTC	ATTTTGGACT	CAAAATTATG	CAGACTTTTT	TTCGGAAAAG	AAAGAAAAAC	11340
AAAATTTTCGA	AACATTTGAA	CCGCGGGAAC	TTCTTTTACA	AATCTCTGAG	GAAGAAGCTG	11400
TCCCGCATAC	AGAAGATTTT	CAGTCAGCCA	TCTCGCCCTC	TATGGGCCAA	ACTTCACTCC	11460
CTGCTCCTTC	TGTGTCAGAA	TACAGTAGCG	TGCCTCGGTC	AGCTTTTTTAC	CCTCTCAGAG	11520
AACGTATCCA	AGAGAGCATT	TCAAAGGCAG	TCATCCCTCC	TTTGACAGGC	TATGTCGGAA	11580
AACAAATAGG	TGAAACTATT	TTCCCTGGTA	GTGGAGATCT	TGTAGCAGCC	GCTGCGTCTT	11640
TAGTTGCAGC	ACAATTGGTT	GATTCAAGGT	TTAATAACAG	AAGACAAAGA	TTGAAAGACG	11700
CAGCCAGAAA	GCGTCACCGC	TATGTTAGAG	AGATGCATAA	TATTTCTGAT	AAAGAGTCAA	11760
ATGCTTCTAA	TGATACGGTA	ATATCACCTT	TGATTGGACA	TGGTTCGGCG	ACTGAAAATC	11820
GTTTTGAATA	TTTGAGACCT	AAAGGTGGAA	ATTATTTATA	CTAATAAAAA	TCATAACAGA	11880
CCTGACGGGC	GGTCATCCTT	TTTTATTAGA	TGCAGAAATT	TGTACCTCCA	CCACGAATCC	11940
TTGCTCCAAC	AGAGGGTAGA	AACAGTATTA	CTTATACGCC	TCTGGCACCA	CTGCAAGATA	12000
CAACAAAAGT	ATTCCTTTAT	GACAATAAGT	CTTCGGACAT	TGAAAGTTTA	AACTTTACTA	12060
ATAATCACAG	TAACTTTTTT	ACAAATATTA	TTCAAATGTC	TGATTTGGCA	GCGGATGAAG	12120
CAGCAACGCA	AGATATTAAA	CTGGATGAAA	GATCTAGATG	GGGCGGTGAA	CTGAAAACCT	12180
TTATAAAAAC	AAATTGCCCC	AATGTTTCAG	AATTTTTTAA	CAGTAATAGC	TTTCTAGCCA	12240
GATTAATGGT	AGATAAAACT	GATCCAGAAC	ATCCTAAATA	CGAATGGGTA	CAAATTACAA	12300
TTCTTGAAGG	CAATTACACT	GGAAGCGAAC	TTATAGATCA	ACTTAACAAT	GGTATTTTAA	12360
ACAATTACTT	AGAAGTGGGA	CGCCAAAAAG	GAGTAGAAAT	TGAAGACATA	GGAGTAAAAAT	12420
TTGATACAAAG	AGATTTTTTCA	CTTGGATATG	ATCCTGAAAC	GGGACTAATT	ACTCCAGGAA	12480
AATATACATA	TAAAGCTTTT	CATCCAGATA	TTATCTTGCT	ACCTGAATGT	GCGGTAGATT	12540
TTACATATTC	TAGAATTAAT	AATATGTTAG	GTATAAGAAA	GAGATTTCCA	TATACTAAAG	12600
GATTTCAAAT	TTTATACAGT	GATTTGACGA	AGGGAAATAT	CTCTCCATTA	CTGAATTTAA	12660
ATAACTATCC	TCATTCTATC	GAACCTGTAA	TGCAAGACGA	AAATGGAGTT	AGCTATAATG	12720

5/23

Fig 1 (cont)

TAGAAAAAAT	AAGTGACAAT	CCCCCAGAT	GGCAAACAAA	GTACAGATCT	TGGACTTTAA	12780
GTTATAAAAA	TAATGGAGGA	GCTAAAGCCC	TAAGTGTACT	AACTGTTCCG	GACATAACAG	12840
GAGGATTAGG	TCAAATTTAT	TGGTCAATGC	CAGATACCTT	TAAAGCACCT	ATTACTTTTA	12900
CTAACAAATAC	TACAAAGCCA	GAAACACTTC	CAATTGTTGG	ATTACATATG	TTTCCTTTAA	12960
AAGCAGGGTT	AGTTTCATAAT	ATAAATGCGG	TTTATTCTCA	ACTTTTGGAA	CAAATTACAA	13020
ATACAACTCA	AGTATTCAAT	AGATTTCTTA	AAAATGCTAT	ACTAATGCAA	CCACCTTACA	13080
GCACCGTAAC	ATGGATAAGT	GAAAATGTCC	CCTTTGTTGC	AGATCACGGG	ATTGAGCCAT	13140
TAAAAAACAG	CCTTACAGGT	GTACAAAAGAG	TTACTATAAC	AGACGACAGA	AGGAGATCTT	13200
GTCCATACAT	ACAGAAATCT	TTGGCGACTG	TTGTCCCCTAA	AGTACTTTCA	AGTGCTACAC	13260
TTGAGTAACA	ATCTGGCTGA	TATCTCTGGG	CCTTATCCTC	CTGGAACCGT	TATGTCTATT	13320
TTAGTTAGTC	CCTCTGATAA	TACCGGGTGG	GGTATTGGAA	CATCAAGTAT	GAGGGCTACT	13380
GGCTTGAAAT	TTTCTAAAAA	ACAACCTGTT	AGAGTGGGAC	CTTATTACAG	AGCTCAGTGG	13440
GGACAGCTTA	ATGCTCGTAC	TTCACTTGAG	AAACTAAAAA	CCAAATTGAA	ATATTATGAA	13500
AAATTGTACA	GGGACAGACT	AAAAAGAAAA	ACAGTTGTTT	CAAGAAAAAA	GAGGTCACCT	13560
ACATCTCCTG	CGGATCGACT	TAAAAAATAT	CTTAAAGCTG	TCAGTCAAAT	CAAAGCTTTT	13620
AATAGAGCTA	GAAGAGCAGC	CCAATAAATA	TTATTTTTC	CTTGAGATG	AAGGTAGTTT	13680
ACGTGCTTAA	ATCTCCTCAT	CGTCGAAGAC	ATACACGTCG	TTACAAAAAA	CTAAAAAA	13740
TCAATCTATC	TCCATACATT	TTACCTAAAG	AATTGCAAGG	CGGTTTTTTA	CCAGCTCTCA	13800
TTCTATCAT	AGCAGCCGCA	ATTAGCGCAG	CCCCTGCTAT	AGCTGGAAC	GTAATAGCTG	13860
CTAAAAATGC	TAATCGTTCT	TAAAATTTAG	AAAACCTTTT	TTTTAACAGA	TCACATGGCT	13920
TTTTCAAGAT	TAGCTCCCCA	TTGCGGCTTA	ACACCTGTTT	ATGGCCACAC	CGTTGGAATC	13980
TGTGATATGA	GAGGAGGTTT	CAGCTGGTCT	AGTTTGGGAA	ATTCTTTTAC	TTCTGGTTTA	14040
AGAAACATAG	GTTCAATTTAT	ATCAAAATCT	GCTCAAAAAA	TAGGTCAATC	ACAAGGATTT	14100
CAGCAAGCCA	AACAAGGTCT	ACTGCAATCA	AATGTTTTAG	AAAATGCAGG	ACAATTAGCA	14160
GGTCAAACTT	TAAATACTTT	GGTAGATATT	GGAAGATTAA	AGGTAGAGAA	AGATCTAGAA	14220
AAATTGAAAC	AAAAAGTTAT	AGGGAACGAC	CAACAAATTA	CTCAAGAACA	ATTAGCTCAA	14280
CTAATAGCCA	GCTTAAAACC	AAAAGATGAA	ATGTTTGTA	AGCAATCAGA	AAAAATTGTT	14340
GAACCTATGA	GACCAGAAAT	TAAATCTAGC	CAAATGCCTG	TAGAAATGTC	TTTTTATGAT	14400
TCTGTAAGTG	ATGAACCAAT	CATAAAAACC	AAAGAAGTTA	GCCCTCCTTC	ATTTTCATCT	14460
GAATCTTCAC	ATTCATATT	TCACCCAAGA	AAAAGAAAAA	GCGTATCCGG	TTGGGGTGCA	14520
TTTTTGGATA	ACATGACTGG	AGATGGAGTA	AATTTTAATA	CAAGAAGATA	TTGTTATTAA	14580
AAACACTTTT	TATTTACAGA	TGGAGCCACA	GCGTGAATTT	TTTCACATTG	CGGGTAGAAA	14640
TGCAAGGGAA	TACTTGCTG	AAAATCTGGT	ACAATTCATC	TCTGCCACTC	AAAGTTTTTT	14700
TAATCTTGGA	GAAAAATTTA	GAGATCCTTT	TGTAGCTCCA	TCGACGGGTG	TAAGTACTGA	14760
CCGTTCTCAG	AACTTCAAC	TTCGTATAGT	TCCGATTCAA	ACTGAGGACA	ATGAAAACCT	14820
TTACAAAAC	AGATTACTT	TAAATGTAGG	AGATAACAGA	GTTGCAGATC	TTGGAAGTGC	14880
ATATTTTGAC	ATTGAAGGAG	TTATTGATAG	AGGACCTACT	TTTAAACCTT	ATGGAGGGAC	14940
AGCTTATAAT	CCATTAGCCC	CAAAATCAGC	TTTTCCCAAT	GCAGCTTTTA	TGCATACTGA	15000
TGAAGCTACA	ACAATTTATA	TTGCTCAACT	CCCTAATGCT	TATAATGCTC	AAAACAAAGG	15060
TGTAGAAGAA	GCAATTCGAG	TAGAAGCAAA	CACTACTACT	CCTAATCCTC	AATCAGGAGA	15120
ATATGCTACT	TATGACTCTG	CCAAATTTAA	TCCAGAAACT	ACTGGTGCTT	CTGGAAGGCT	15180
TTAGGAATT	AATAGCTTAG	GAGATCTTTT	TCCGGCTTAT	GGATCTTATT	GTAGACCTCA	15240
ATCAGCAGAT	GGTAACATTT	CAACTGCACC	CATAACTAAA	GTCTATCTAA	ACACTACTGC	15300
TACAGATGAC	AGGGTCAGTG	GAGTTACTGC	AGTTGACACC	GCAACCAGAT	TGCATCCAGA	15360
TGCTCATTAT	ATTGAATATA	CTGATGAAGC	CAAAGCTACA	GCTATAGGAA	ATCGCCCAAA	15420
TTATATTGGT	TTCCGAGACA	ATTTTATTGG	ACTCATGTTC	TACAATAATG	GTTCTAATGC	15480
AGGAACATTT	TCCAGCCAAA	CACAACAAT	TAATGTTGTT	TTAGACTTGA	ATGACAGAAA	15540
CAGTGAAC	AGCTATCAAT	ATCTAATAGC	AGATCTGACA	GATAGGTATA	GATATTTTGC	15600
ACTTTGGAAC	CAAGCAGTTG	ATAGTTACGA	CCAGTATGTC	AGAATTTTGC	ATAATGAAGG	15660
ATATGAAGAA	GCCCCCTCCG	CCTTATCATT	TCCTTCTCAA	GGTATCCAAA	ATTATTTTCAT	15720
GCCTACTGCG	GCAGGTAATG	CGATGACAGT	AGACACGGGT	AGAAATACTG	CAGCAAAAAC	15780
AGATAACACC	AAGGCTTTTA	TAGGATATGG	CAACATGCCA	TCTTTGGAAA	TGAATCTGAC	15840
AGCAAACTCA	CAACGTACAT	TTTTGTGGTC	TAATGTAGCA	ATGTATCTGC	CAGATAGGCT	15900

6/23

Fig 1 (cont)

GAAAAACAACA	CCACCCAACA	TAAATCTACC	TGATGACACC	AACTCTTACG	GATATATAAA	15960
TGGAAGGGTC	CCTCTAGCAA	ACATAATAGA	TACATGGACT	AACATTGGGG	CTAGGTGGTC	16020
ATTAGATGTT	ATGGATACTG	TAAATCCATT	TAATCACCAC	AGAAATTCAG	GACTAAAGTA	16080
TAGGTCACAA	CTGTTAGGAA	ATGGAAGATA	TTGCAGATTI	CACATTCAAG	TACCTCAAAA	16140
ATTTTTTCTT	ATAAAAAATC	TTTTGTTGCT	GCCAGGAACA	TATAATTATG	AATGGTACTT	16200
TAGAAAAGGAT	CCCAACATGG	TTTTTCAGTC	TACTTTAGGT	AACGACCTTA	GAGCAGATGG	16260
CGCAACTATT	ACATACACCA	ACATAAATTT	ATATGTTTCA	TTTTTCCCTA	TGAATTATGA	16320
AACAGTAAAGT	GAACCTTGAAT	TGATGTTGCG	TAATGCTACT	AATGATCAAA	ACTTTGCAGA	16380
TTATTTGGGT	GCGGTAACCTA	ATCTTTATCA	AATCCCAGCT	AATACAAATA	CTGTAGTAGT	16440
GAACGTACCA	GATAGATCTT	GGGGTGCTTT	CAGAGGATGG	AGTTTCAATA	GAATTAAAGC	16500
TTCAGAAACA	CCTATGATAG	GAGCAACAAA	AGATCCAAAT	TTACTTATT	CAGGATCTAT	16560
ACCGCTACTA	GATGGTACTT	TCTATTTAAC	ACACACTTTT	CAACGAGTTT	CTATTTCAGTG	16620
GGATTCTAGC	GTTCCATGGC	CAGGAGATGA	TAGGCTTTTG	ATTCCAAATT	GGTTTGAAT	16680
TAAGAGAGAT	CCTAATATGG	ACGCAGAAGG	TTATACTATG	AGTCAAAGTA	CTATCACAAA	16740
AGATTTTTAT	TTGGTACAAA	TGGCTGCTAA	TTATAATCAA	GCTTATCAAG	GTTATAAATT	16800
GCCAGTACAT	TCTAAATATT	ATGGATTTTT	AGAAAATTTT	CAACCTATGA	GTCGCCAAGT	16860
ACCAATTTAT	GGTAATGGCA	CTTATGATTT	ATATACTGCT	TATATTACAA	ACCAAAGAAC	16920
CATGCAAATT	TGGAATAATA	GTGGTTTAGA	ATCTAAAAC	TCAAATCCTC	CTATGTTATC	16980
CAACACTGGT	CATCTTTATG	TAGCTAACTG	GCCATACCCT	TTGATTGGAC	CAAATGCTAT	17040
TGAAAACCAA	CAAACTGAAA	GGAAATTTTT	GTGTGATAAG	TATATGTGGC	AGATACCATT	17100
TTCTAGTAAT	TTTTTGAATA	TGGGTAATTT	AACAGATTTA	GGGCAAAGTG	TTTTGTACAC	17160
TAATTCCTAGT	CATTCACTTA	ATATGGTTTT	TACTGTGGAT	AGTATGCCCTG	AAACAACCTTA	17220
TCTAATGCTT	TTATTTGGTG	TTTTCGACCA	AGTTGTTATT	AATCAACCAA	CAAGAAGTGG	17280
AATAAGTGTA	GCTTATTTGC	GCCTTCCTTT	TTCAGCTGGT	AGTGCAGCAA	CATGAGCGGC	17340
ACATCCGAAA	GTGAGCTGAA	AAATCTGATT	TCATCATTAC	ATTTAAATAA	TGGATTTTTTG	17400
GGCATTTTTTG	ATTGCAGATT	TCCAGGTTTT	CTGCAAAAAT	CTAAAATTCA	AACTGCTATT	17460
ATTAATACAG	GTCCAGAGA	ACAAGGCGGA	ATACACTGGA	TAACATTAGC	ATTAGAACCC	17520
ATTTCTTATA	AGCTATTTAT	ATTTGATCCA	CTCGGATGGA	AAGACACTCA	ATTAATTAAA	17580
TTTTATAAAT	TTTCACTAAA	TTCTCTTATT	AAAAGGTCGG	CCTTAAATAA	CTCAGACAGA	17640
TGTATTACAG	TAGAAAGAAA	TACTCAAAGT	GTTCAATGTA	CCTGTGCGGG	ATCGTGCGGC	17700
TTGTTTTGTA	TATTTTCTT	ATACTGTTTT	CACTTTTATA	AACAAAATGT	ATTTAAAAGT	17760
TGGCTTTTTTC	AAAAATTAAA	CGGTTCAACC	CCTTCTCTGA	TCCCATGTGA	ACCACATCTA	17820
TTACATGAAA	ACCAGACATT	TCTTTATGAT	TTTTTAAATG	CAAAAAGTGT	TTATTTTCGA	17880
AAAAATTATA	GAACATTTAT	TGAAAATACT	AAGACTGGAT	TAATAAAAAAC	ACATTAATTG	17940
TATTCCTTGCT	TTTTGACGTT	TTTATTAGTC	TTCATCTTCA	TCTTCTTCTT	CACTGCTAGA	18000
TTCCAAGATG	GTTTTTTTTT	TCTTTGATGG	AGTAGGCTCT	TCAATAGTTC	CAAAAGGATT	18060
CATATCAGAA	TCCTCTTCTA	TGTTAGGCAA	CATAGTATTT	TTAACCTGGA	ATGACTGATT	18120
CCACTTAAAT	TGAGAAAAC	GAATTGGAAT	GTTATTTCCC	ATACATTCAT	TCCAAAATTT	18180
ACGCACAAGA	GTTAAACACT	GTAACATATC	TGGCAAGCTA	ATTTTCATCT	CACAAAATTT	18240
TCCATTATTA	CGTCTCAAGT	TGTATTGATA	GTTACAACAT	TGAAACACAA	AAACAGCAGG	18300
GAATGTAACT	GCTGCGGCC	GAACCTTATT	AACATCCTGA	ACATCAATTC	CTTCCACTCC	18360
AGATATAGAA	AATGGAGTTA	TTTTAGGGAG	TTGTTTTCTT	ATTGTTTGTT	TGCCACCATA	18420
ATTACATTCA	CACTGACCCA	ATATAAAAAG	CATATTTCCG	ACTTTAGCTT	TCGGAACAC	18480
AGCTTTTGTA	GTTTCAATGG	CATTTTGCAT	AGCCAGCAAG	GCCTTCTTTT	CATCTGAAAA	18540
GTTAAGACCA	CAACTGCGAG	GAGAACATTG	CCCAAAACGC	TGATGGGCAT	CCTCAGCACA	18600
TAACACGTAA	TGTTCCCTGAA	CTATTTTTAC	TACTTGTTTA	TTCATACGCC	CATTACTAAG	18660
AACACCCCTC	CCTTCCCTTA	GGGCTTGCAC	CCCTGCTTCC	GATGTTGGAG	GCATTTCAAT	18720
TTCAATCACC	CTTTTAAACA	TGAAGTCACC	ATGAAACAT	CTAGGACGGT	CCTCCTCCCA	18780
ATCATGATAC	CACAAATAAC	AACCAGAAGC	ATTAAAGTTT	GGAATCAAGT	CAATTTGCTT	18840
ACAAATTGCA	CTATATAGCA	TTCTACCTCC	TACAGTAGCC	ATAGATTTAC	TGCTACTATA	18900
AGTCAAATTT	ATAATTTTCA	TCTTTTTTCA	GTAAGTACCA	AATAATTTTT	CACAATCTCC	18960
TTCTTCAGGA	TGAAACTTCA	TTTGACTGGT	ATCAACTTTA	ACACACTCTC	CAATTTAGC	19020
TAAATTTTCG	AGCGCCGCTT	GAACCTTATT	CTGAAATCTT	TCTGTAGTAG	ATTTTCTCTT	19080

7/23

Fig 1 (cont)

CTTGATAGAT	TTAGTAACTT	TTTTAGAAGA	CATTATGTTA	GTTTTTTTCT	CGTTGTAGGA	19140
TGGCTGAAAA	AAATATGGGA	GAGTCAGAGA	AGGGTTTGAA	CGAAGAAGAA	TTTAACTCTA	19200
TTCTATCAAA	ACATCTGGAA	AGACAAATTA	AAATCTGTAA	AGCGTTAACA	TCAAAATTAT	19260
CGAACTGGAA	TATTGGAACA	TTGTTAGAAA	ACTTGTATT	TTGTCCTGAT	GAAAGACAAT	19320
CATCAGGTGA	TCCCGACCCA	AAACTAAACT	TTTATCCGCC	TTTTTTAATT	CCGGAATGTC	19380
TTGCATTGCA	CTATCCATTT	TTTCTAACAA	CTCCTATTCC	GCTATCATGC	AAAGCGAACA	19440
AAATAGGAAC	TAACACTTAC	CGAAAATGGA	TGAACAATCA	AGTCCTGGAT	TTACAAATAC	19500
CTTCCTTGGA	AAATTGCAAA	TGGGATGATA	GCTTGGGAAA	TGTAGATTTA	ATTGAAGAGC	19560
TTAAAGAGAA	CCAAAAACTT	GTTTTAGTAA	AACAAGACCA	TGAAAGAAAT	ATATGGTTTA	19620
AATCAAAATG	CAAACAACCT	CAAAGTTTCA	GCTATCCCTC	ACTCAGTCTG	CCCCCAGTTT	19680
TACAACAAGT	TTTAATTGAA	TCTCTTATCG	GCATTAGTCA	GGATCCTAAT	AACTTTGACA	19740
AAAATTACGA	ACCTGCAATA	ACTCTAGAAA	AACTACAACA	TGTAAACTGT	GATCAAGATT	19800
TAAAAACAAGT	TCAACAAAAA	GTATCTTCAG	CCGTACATA	CGGAATACTT	TTGAAATGCA	19860
TTCAGACTTTT	ATTCACTGAC	AAATTATTCA	TTCAAAACTG	CCAGGAATCA	TTACATTACA	19920
CCTTTAACCA	TGGTTATGTA	AAATTACTTC	AATTTTGTAC	AAATGTCAGT	TTAAGCGAAT	19980
TTGTAACTTT	CCATGGTTTA	ACACACAGGA	ACAGACTCAA	TAATCCGCAG	CAACATACAC	20040
AATTGGCAAC	CGAAGACAAA	ATAGACTATA	TCATAGATAC	AGTGTATTTA	TTTTTGGTAT	20100
TTACGTGGCA	GACAGCAATG	GATATTTGGA	ATCAAAACATT	AGATGATAAA	ACAATAAATA	20160
TAATTAAGA	GGAATTA AAC	CAAAATTTTG	AGAAAATTGT	CAAAGCTGAA	TCAGTTGATG	20220
AAGTTTCTGA	AATTTTAAAG	TCTATTATTT	TCCCTGAAC	CATGCTGCGA	GCTTTTTGTT	20280
CTAATTTACC	TGATTTTATA	AATCAGAGTC	AGATATCAAA	TTTTAGAAAC	TTTATCTGCA	20340
TTAAATCCGG	CATACCGCAG	TCAATTTGCC	CCCTATTACC	TTCAGATCTA	ATTCCTTTAA	20400
CTTTCCTAGA	AAGTCATCCA	ATACTCTGGA	GTCATGTAAT	GTTACTAAAT	CTTGCTTCAT	20460
TTCTAGTAAA	CCAAGGCAAT	TATTTGCATG	AACCCGAAAA	ACCTTTAAAT	ATTTTCATCAG	20520
TTTACTGTAA	TTGTAATTTA	TGCTCTCCGC	AAAGAATGCC	ATGTTACAAT	AGCAGTTTGA	20580
TGCAAGAAAT	ACTAACCATT	GATAAATTCG	AGTTCACAAA	CTCTGATAAA	ACAAAACAGC	20640
TAAAACCTGAC	CCTCCAAACT	TTTGCTAATG	CCTATCTTAA	CAAAATTTAAC	TCAGCAGAAAT	20700
TCTAACCATGA	CCAAGTTTTA	TTCTACAAAA	ACTGTAAAAG	TAAATTTTCT	AACCAATTAA	20760
CAGCTTGTGT	AATAAAAGAC	GAAAAATTAT	TGGCTAAAAT	AGCAGAAATT	CAAATAACGC	20820
GGGAAAAAGA	ACTCTTAAAA	AGAGGAAAAG	GAATTTATTT	GGATCCAGAA	ACAGGAGAAA	20880
TCTTAAACAA	TGGAGAAGCC	ATATCATCCT	CTGAAAACCT	CCAAAGGCAA	AGAACTAGCT	20940
ATGCTCTACC	ATCAAAATGAA	GGAGAGCGAG	CTGGATGGGA	AGCCGATGAG	CGAAGAAGAC	21000
GAAGGAGAAG	TGAGTGAGGA	TGAAACAGAG	ACAACAATTC	CAAAGAAAAT	GAAGTTTACA	21060
AGTAAGTAAG	CTCTAAATTT	TTTATATTAA	AAACTGAATT	TTTTTAGACA	AAATTATTTT	21120
AAATTAATATC	TTTATAGCTA	GCAGTTGATC	TTTGTTCGTT	TTTCAGAAAA	CTCAAGTGTT	21180
CAGTCATATC	AAGTTCACCT	GCCTCTGAAA	CACGAAATTC	CGGAAATTC	AGAAAAAATT	21240
AGACTAGAAT	CTAAAAAATA	TCCAGGAAAA	GTTTATCAAA	TAAGAAATAG	AACTCCAGCA	21300
AGTATTACAA	AACGATACCT	GTATGAAAGA	GATCTGAAGA	AACTGTTCCA	GTATCTAGAA	21360
GACGCAAAAG	AGCTTTACGC	TAAGTACCAA	AGCTGAGGCT	TTATAGTTTT	AAATTTTCCC	21420
GCCATGGCTC	AACCACTGAC	GCCTTACGTC	TGGAATATACC	AACCAGAAAC	AGGATATACT	21480
GCTGGAGCCC	ATCAAAATTA	TAACACTGTT	ATCAACTGGT	TGCATGCCAA	TCCACAAATG	21540
TTTGCCAGAA	TTCAACATAT	AAACACCGCA	CGCAATGTTA	TGGACAAATT	CCGCTCTGAT	21600
TTGACCCGAG	ATGACATCGC	GGTTAACATC	AACAACCTGGC	CTGCAGAGGA	TTTAATGCAA	21660
CCTCCTAATT	TTCTTTACAT	TCCTGCGACC	TCTAAATCCG	CTTCAACCAT	AAATGACTGG	21720
TTGGCTACCA	CTCAAGGAAT	TCAACTCAGT	GGAACCTAGT	AACTAAACGG	GTGGGGATCT	21780
AACCGCCTGA	CTTCCTATCC	GGATATTCCA	CCCATTTTAA	AGTATGAAAG	GCCTGGTCAA	21840
CAACTTCAAG	GCCAAGGACT	TTTTAAGCAA	GAAAATATT	ATTTATTTTA	CGAATCTCCG	21900
CGCCTCCCTC	GCTCTGGAGG	ATTAACTCCC	CAACAATTTG	TAAAAGAATT	TCCGCTGT	21960
GTTTATAATA	ACCCCTTCTC	AGAATCTATG	AGTGTATTTT	CGAAAGAATT	TAGTCTTTTG	22020
TTTAACCCTT	CAGAATCTTT	GAAAAAACA	TCCAGTCAAA	CTTTACAATA	TAAATAAAAA	22080
ACTTCTATTG	ATCTTTATAC	TTACACTAAA	GCATCGCGTT	TATTTTCGTC	GCCATAAAAA	22140
TATATCAAAG	ACCCGTAATT	CTCTAACTTT	AAATCATTTT	TTGAACTAAT	CTTAATCCAT	22200
TTAAATGTAG	GAATTAATAT	ATCAGAAACC	AGTAACAAGC	CAGAATTAAT	ATATACTTGT	22260

8/23

Fig 1 (cont)

GTCATTTTTA CAGATGAAGC GAGCACGCTG GGACCCGGTT TATCCCTTTT CTGAAGAGAG 22320  
ACTGGTTCCT CTGCCTCCTT TTATTGAAGC CGGAAAAGGG CTAAAAAGCG AAGGGTTGAT 22380  
CTTATCTTTA AACTTTACTG ATCCTATCAC TATAAATCAA ACCGGTTTCT TAACGTGAAA 22440  
ATTGGGAGAT GGAATATTCA TAAACGGAGA GGGTGGCCTA TCAAGCACTG CTCCAAAAGT 22500  
CAAAGTTCCC CTGACTGTCT CAGATGAAAC ATTGCAACTG CTATTAACTA ATTCTCTAAC 22560  
AACTGAGTCA GACTCTTTAG CTTTAAAACA ACCGCAACTT CCCCTAAAAA TAAATGATGA 22620  
GGGGAGTTTA GTATTGAACT TAAATACTCC TTTAAATCTA CAAAATGAGA GATTGAGTTT 22680  
AAATGTTTCA AATCCACTAA AGATAGCGGC AGATTCTTTA ACTATAAACT TAAAGGAACC 22740  
CCTAGGATTG CAAAATGAAA GTTTGGGCTT AAATCTAAGT GATCCTATGA ATATAACTCC 22800  
AGAAGGAAAT TTAGGTATTA AATTGAAAAA TCCTATGAAA GTTGAAGAAA GTTCTTTAGC 22860  
CTTAAACTAT AAGAATCCTC TCGCCATTAG TAATGATGCG TTAAGTATAA ACATTGCGAA 22920  
TCCATTAACAT GTTAATACAA GCGGATCTCT AGGAAATATCT TATTCTACTC CCTTACGAAT 22980  
TTCAAATAAT GCTTTATCAT TATTTATAGG AAAACCTTTA GGATTAGGAA CTGACGGCTC 23040  
TTTAACTGTA AATTTAACTA GGCCTCTGGT ATGTCGTCAG AACACTTTGG CCATAAACTA 23100  
CTCAGCCCCA CTAGTGTCAT TGCAAGACAA TCTTACTTTA AGTTATGCTC AACCATTAA 23160  
TGTAAGCGAT AATTCTTTAA GATTGTCTCT AAATCTCCA CTAAACACAA ATAGTGATGG 23220  
AAAACCTTAGT GTAAACTATT CTAATCCTTT AGTTGTGACT GACTCTAATC TTACCCTCAG 23280  
TGTTAAAAA CCTGTAATGA TTAACAACAC AGGTAATGTT GACTTAAGCT TTACAGCTCC 23340  
CATAAAATTA AATGATGCAG AACAGTTGAC TTTAGAAACC ACTGAGCCCT TGGAAAGTGGC 23400  
CGATAACGCT CTAAAACTGA AACTTGGAAG AGGCTTAACT GTTAGTAATA ATGCTTTAAC 23460  
CTTAAACCTT GGAAACGGTT TGACTTTCCA ACAAGGTCTT TTACAAATTA AAACCTAATAG 23520  
CTCTCTAGGG TTTAATGCTT CTGGGGAATT ATCAACAGCT ACAAAGCAGG GAACCATAAC 23580  
CGTTAACTTT CTAAGCACAA CTCCTATAGC TTTTGGGTGG CAAATAATAC CTACTACTGT 23640  
AGCTTTCATT TATATTTTAT CAGGAACACA ATTTACTCCT CAATCCCCAG TAACCTCTTT 23700  
AGGTTTTCAA CCCCACAAAG ACTTTTTGGA TTTCTTCGTT TTAAGTCCGT TTGTTACATC 23760  
TGTAACCTCAA ATTGTGGGAA ATGATGTAA GGTATTGGC CTAATATTT CTAAAAACCA 23820  
ATCTACCATA ACTATGAAAT TTACTTCTCC CTTAGCTGAA AATGTACCAG TTAGTATGTT 23880  
TACAGCAT CAATTCAGAC AATGAATATT TTAATAATTC TTTATTAAG AGTAATCTTT 23940  
TTACATACCG TTCTTGACAT AATGTGCTC TATAATTAAC AAATCTAAGC AAGCAAGGTT 24000  
GATCATTGGA ATCTATAGAA GCATAACTCT TCCAATAAGC ATAATCATAT GGCGGTAAAT 24060  
GAAAACCCCT TAAATCTACC ATATTCATCT TTAAGTGATC AGTATCTAAC AGGTTTTTAC 24120  
AATCTTGAC TTTCTGGACT TTAATAACAA ACAGTACTTT CATAGGACAA CAATTGTAAC 24180  
GGTTATAATC TGTTACAATT TTACTTATTT CTCTTCCAA TGGCAAAGCA TTCCAAAGTC 24240  
TTGTTATAAG TACTGTAAAA TCATCAAATG AATAACATAA CACATTTGTA CAACAATTGG 24300  
TCCAAGGTAA AAAACAGGC ACACGAACAT GAACTTTTTT TAAATTAAC ATCAGTGTCT 24360  
GTTTTAAACT TTGACATTGC AAAGAATTTG GCTGCAAGCA ATGACAATGA AATTGATTTT 24420  
GCTGACAAGG TAAGTCACAC AAATACAAC TTAACAGCCT AAATATAACA ACATTAATGT 24480  
AACTTTCCAA GACTTTAAAA CTAACAAACG GTATATCACA ATAAAAAGA TGATGAATCC 24540  
CTTCGCAACA CATAATGGAG TTCATGCTAC ATCCAAAGAT GGTTCCGACA AACCTCTGTA 24600  
AATTAAGAA CAACAATACA ACATACGAAG AAAATTAATA CGTTTTTCAA AACGAGATAT 24660  
ACATTGCTGC AAAGTATCTG AACATTTACA TTTTATACTT ATAAGCTCAC AAGTTTCAGA 24720  
AAATGTAATT CGTTTAACAG TTTGATATGA ATACCATTTT GAAGAAAAAT AGAAATAGTT 24780  
TTGTGCATTT GTAAGCTCCC AGAAACATTA ACGGACAGGC AAATCCAAGT ATTACAACAA 24840  
ACAGGAACAG TCTTAACGTT TCGTTCAGAA AACAAAGTAA CAGGCATATG ATTAAGCAA 24900  
GACAATAAAA CACTTTTGGC AGCTAAACAT TGCAAAGATC CAGGTGAATT ACAATGACAA 24960  
TGATAATAAAA ACTTATAAGC CATATCGGCC CTCTTGCAA ACGAATCAGC TTTTGGCTT 25020  
ATAGGAAAAA AACAAAAAAA CTGATTATAT ATGAATGGAG TTAATATCTT CTTCAAAATTA 25080  
TACACACGAA TAGCAGAAC AAGACGACCA CGCCCAACAC AGGTAAATAT TTCAAGTCCA 25140  
TGACTAGGAA CAGATGGTTT CTCACAAGCA ACACTTTGA TTTGCTTATC CATCACTGCC 25200  
AATCAGGCTT AATAGGAAAA GAAGAAAAAT AATTTTCCA ATAATAACGA AAGAAATTCC 25260  
ACGTTTCATC CTGTACATTA CTAGTCACAA ATACAACCTC CGCTATCAA GATTCCCTAT 25320  
CATTTAAAC TCCCACCAA TTGTCCAGT CTACCTCAA AAAGCCAGTT CCCATATTTT 25380  
CAAAATTTGC CCATTTTAAA TAATCCAAAG CATCAAATTC AGGAAACAAA TCTTCTGAG 25440

9/23

Fig 1 (cont)

CTAAACATA	TACAGTTT	TCGCCATTA	ATCTAAAGC	CATCCTAAAT	GGACCTCTAG	25500
CCCAGTAGTT	TAAGTACCGG	GAAGAGACTA	TACAATATAC	TTGATATTGA	TGTCTGTAA	25560
GTGGTGATAA	AAAAGAAAGT	AATTCAGAA	TAGGATAAAG	CATTCTCCCA	TGTTGATTCA	25620
TCTACAAAA	ACAAAAAAT	TATAAGCTTC	ATAGAAAAAC	TACTATTTAA	CAAATCTATA	25680
AAAATGCATT	AAAAAGTTAC	CTTGAATATA	AATTCAGATC	ACCTAAAAAA	CGAAAAAAA	25740
TAACATTTAT	GTTAGTAAAT	GATAGTCTTT	AAAAATTAGA	AAAGAATCAA	GTCGCTTTTA	25800
TACTTACAAA	CTCCAAATAA	ATTCTGTAA	CAAGAGAAAA	ATTGTAACCT	AAAAGGTAAA	25860
GAAGAACATT	ATAAGATTAA	AACCACTCTA	AAATCTGAAA	AGCATTATGA	AAAATTCTGA	25920
TAGCTGCAAC	TTACTAGTCT	TCTCCAAATG	TTGCAGGCAT	TTCAAAAAAT	CAAGAGGAAA	25980
ACCGGAGTTT	ATAAGTAGT	AGTCTGATTA	TATCTGAAAA	AGTTTAACTT	CCTTTTCAAC	26040
CCAACCCAGT	CCAATAAAAT	TCCAACCTTA	ACTTCTTTCC	TGCTAAAACT	CCATAAAAGT	26100
CCAATTACCA	CTTGACTTTT	ATTTAACCTC	AATTATGTTA	CATGTTATTC	TACCCATAAA	26160
AACTTGATGA	CCAAGAAGT	ACCTTTCCCA	TGTTTTCTG	AAATAACAAA	AATGTTGATT	26220
TAAGATTTT	TAAGTACCCA	AAAAACCCGC	TCTCATGATT	TTTTCTTATA	TAAACAGGAT	26280
ACAAAAGAAC	TGCAAAAGAT	ATTCCATCAT	ACTTCTCCAA	CTGTCAAAAC	ATACCACTTA	26340
ACCTCTCCCA	TGTTTTTTCC	CTTTTGCACA	AACAGGATAT	AAAAAATATT	TTTGCCACAA	26400
TGTTTTTCTT	TTTACTCAAC	TGCCAGAATA	AAAATGAACA	GCTTAACCTT	TTTCCCTCTT	26460
AACCCATTGC	GTTCTCTTAA	GAAAAAAATT	ATCCCGCCCA	ATATGCTAAA	GGCTTCTCCC	26520
GCCAAACAG	CTCAACTTAA	AATCTCTCAT	GAATAAAACC	CAGAGAAAA	TTCCAGTAAT	26580
AAAAATTAAT	AACCGTGAAG	TACTAGATCT	AATAATGATA	TTTTGAACCT	ATAAAAAATCC	26640
ACCATCCATG	TAATGTTACA	AACACTTTTT	TATTGAGTTT	TTTCTTACAA	CTGCATTACA	26700
TACAGGCCAA	GCATCAAACT	TTCTTCTGTA	TTTCTTCTTA	GACCACAAAA	TTACAGACTT	26760
ATATTTCTGC	CACAAATCTC	TATGATCTTT	ACAGTAACAC	TTACATTTAA	ATGGGGAATA	26820
CAGCAGCAAA	TAAGGATGAG	TTAAACATGC	GATACAATGA	CCAGAAGGAA	GATAATACAA	26880
TACATCACAC	CAAAATGAAG	GTACAGACAA	CATCGCATGA	AATCTTAAAT	GTGATTTTAC	26940
AATAAATTTT	TGCAGCAGCT	TACAATCTAT	ATTAGCAAA	CGTTTTATAT	ACAAACATAA	27000
AACTTTGGAA	CTTTTCACCA	ACTCAATCAT	GTTATTATAA	CACATTACAA	ATTTTGCTAT	27060
ATCTTTATTT	GTCAAAATAAC	AAAATATCTC	AATCCACAGC	TCATCTGGCA	GCAAACCTCG	27120
CAAAATCCATG	ACCTGTAAAA	GATACAACAG	AAAACAGAAA	ATTAATGCCA	TTCAATAACA	27180
TAAAAAATAC	AGTCAAAATCA	CATACTTTTT	CTCACTTACA	AAACTTTGTG	AGCAGGCCTC	27240
CAAAACAAAC	TTACAGAAAT	GGATGCATAC	AAGAACATTC	TCCTCTCAAA	AATTGCTTTA	27300
ACTGAATGCG	GCATTTTGCA	CCTCCAGAAA	AATGCAGTCC	ATTGAGAGGC	TCTTCTCTTA	27360
AAACACAGAA	ATGCTTCTGC	AAAATCTGTA	AAGAACTAA	CAACTTCCAA	ATTCCAATCA	27420
TCATGCATTG	CAAAGAAGGA	CATTCAACAG	CAAAAGGATC	GTGATGAGCC	AATAAAGCTT	27480
TACTGTATGA	CTCATTTTCA	TGAATTACAG	TCTGTAACCT	ACTATAATGC	ATTTTAAGCT	27540
CTGCTTCACA	AATTAATAAT	GCTAATTTCT	TTAAGCAGCT	CAAAGAAAAC	TCATCAGGAC	27600
AACGGCATT	AAGAAAGCAA	CAAAATGATT	TCTTAAAAATA	CATTTTTCCT	GCATGATGAA	27660
CAATAAAAAA	TTTCAACGTT	AAACAATGCA	AAAATGCATT	TTTATGCACA	GTGAAAGTAA	27720
TTTTTTCAGC	TGAAGCTAAA	TCACAGCCTA	TTTTATTACA	TGATTTTGTA	TGCTCCAAAA	27780
GAGCTTGTTT	TAATTGCTTC	AAATCCATCT	TCTTACAATT	TTTTCTTTTT	ATAAACACCA	27840
GAACCGCATT	CAGGCCAATT	CCAGTTATTG	TTTAAATTTG	CTACAGAAAC	TGCAGACCAC	27900
AAAACCACAT	CCTCTAAATC	AACCCACAAA	GATCTATGAT	CCACACAAAA	ACACAAAGAA	27960
TGATACGGAG	AATACAACAA	TAAATGGGGA	TTAACAAGGG	ACGCAACACA	ATGACCCGAA	28020
GGTAATAAAG	TTTTACAGCA	CCAATTACAA	GCAACAGGTA	ATGGAGTATA	TTTCCCAATG	28080
CGACGAGAAA	GCCGAATGTC	ATTCAGAACA	GCATTGCATT	TTATCTTCTC	AAACCTCTTA	28140
AGGTGCAATT	GTATAAAATA	AGAATCCTTA	ATGACAGTGA	TGAATTGAGG	AAAAGCAAAA	28200
ACAAAACCTAG	CAATGTCTTT	GCTTGTAAGT	TTCAAAAATA	TCTTCATCCA	AATCTCAGTC	28260
GGTAATTCAA	CAAAAAATTC	AGGGGCCTAC	AAAATTAATC	AGACTAATTT	AATATCATCT	28320
TGTAAACAGC	GAAAAGAAAA	AATAACACAC	CCAAAAATAA	AAAACCTCTTA	CCCCTGTTAT	28380
CCATCGAGAT	ACACAGAAAA	ATTGAGAACA	CTCAGTGTCA	TGTTTCTTAA	ATTGTTCCCA	28440
AAGCTCAGAC	ATTCTAAGCC	AAAAATTTTT	TGAGAAGTGC	AAAAACCCAG	TTTTTATAAC	28500
AAAGCCTTAA	TGTTTTCTTA	ACTGATTTAA	CTGCCCTAAC	AGGAACCTCA	CATTCCGGCC	28560
ACCGCCACCC	AGGGGACAAA	TCTTGCCAAG	AACTACAAGT	CCATAAAACA	ACATCCTGCA	28620

10/23

Fig 1 (cont)

AATTATACCA	AAGGTTTCTA	TGGTCGACAC	AATTACAACC	TGACCTAAAA	GGTGAATAAA	28680
GCAGTAAATA	AGGATGAGTT	AAACAGGCCA	CACAATGTCC	AGAATGTAAA	AAATGCTTTG	28740
TTTGGCACCA	ACCAGACCAC	AGCTGAAGCA	AAGGAAAATT	GTAGCGAACA	CATTCTTCTC	28800
GTAATCTGTT	TAACACAGAA	CAACATTCAA	TTCTGGCAAA	CCTCTTTAAA	AAATGTTTTT	28860
TGAAATATTT	CTTTAAATG	ACAGTTTGCA	ACTCTGGAAA	ACACAAAATA	AAAGCCGCAA	28920
TATCTCTACT	GCTTAAATAT	AAAAATATCA	TTGTCCAAAT	TTCTACTGGT	AAAACGTAAA	28980
GCATCTTCTT	CCTATTAAAA	AAAGAAAAGT	GTTTTCAAAT	TATATTAGAC	TCTAACCAAA	29040
AAAATTCAAA	TACTTTTCCT	TTATAATGTA	CATTAAGAAT	AAAAATATAC	TCACCGTTTA	29100
AAAGTAGAAC	TTAACAGTAT	AATATAAATA	CAAGTGAGCT	GAACAACGAC	AGCCGATTTC	29160
AGCCGGAGCA	AAATTAAAAA	GAATAAAAGG	ATCAAACCAA	CACGTAGGAC	AGTCTACTCC	29220
AAAACAGTAA	CGGCAGTATG	ACACAGAAGG	AGAGGAACTA	AGTCCAGGAA	ACTTCGCCCC	29280
GTGCGATAAA	AAGTAACGCC	GCCGGAAAGC	AGTTGAATAC	AAAAGAGGTA	AAAATTCACG	29340
AAAAACAGAA	GCAAAAACTA	CTAAATCTGC	TATTGGCAAA	TAAAGAAAAA	TTTCAAACCA	29400
TATTTCCAAA	GGAAGAAAAAG	CAATCATACC	GTAGAAGAAC	CTGAAGGCCA	CCGCAAACGT	29460
GCTCCCGTAC	CACAACGTCA	CACGCCACAC	CCACTGGGAA	AACCCACACG	CCCCGCCTCT	29520
GTGCAACGTT	ATATATATGA	ATAG				29544



11/23




Fig. 2

12/23




Fig. 3

13/23




Fig. 4

14/23




Fig. 5

15/23




Fig. 6

16/23




Fig. 7

17/23




Fig. 8

18/23




Fig. 9





20/23





FIGURE 11A


21/23



Infection with OAY204 (MLP/YP7sc)

FIGURE 11B

22/23



OAY206 (HCMV/VP7sc)

FIGURE 12A

23/23

48hr pi




FIGURE 12B

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/AU 95/00453

## A. CLASSIFICATION OF SUBJECT MATTER

Int Cl<sup>6</sup>: C12N 15/34, 7/01, 15/86; A61K 48/00, 39/235

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC<sup>6</sup>: C12N 15/34, 7/01, 15/86

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

AU: IPC as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DERWENT: ovine, sheep, ewe, adenovirus, adeno (w) virus, vector, trichostrongylus or colutoriformis or taenia or ovis or lucilla or cuprina

CHEMICAL ABSTRACTS: STN sequence search

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	VETERINARY MICROBIOLOGY, Volume 41, No. 3 (1994), BOYLE, D B et al. "Characterization of Australian ovine adenovirus isolates", pages 281-291 whole document	1-23
A	NICHOLSON, B H "Synthetic Vaccines" published 1994 by Blackwell Scientific Publications (Oxford, UK) pages 346-361 whole document	1-23
A	JOURNAL OF APPLIED PHYSIOLOGY, Volume 76, No. 6 (1994) LEMARCHAND, P, et al. "In vivo adenovirus-mediated gene transfer to lungs via pulmonary artery", pages 2840-2845 whole document	1-23

☐ Further documents are listed in the continuation of Box C

☐ See patent family annex

<p>* Special categories of cited documents:</p>	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier document but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  
26 October 1995

Date of mailing of the international search report  
6 NOVEMBER 1995

Name and mailing address of the ISA/AU  
AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION  
PO BOX 200  
WODEN ACT 2606  
AUSTRALIA Facsimile No.: (06) 285 3929

Authorized officer

JENNIFER POTTER

Telephone No.: (06) 283 2247